

26 Rec'd PCT/PTO 30 JAN 2001

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE		ATTORNEY'S DOCKET NUMBER <b>EISENBACH 3</b>
<b>TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371</b>		U.S. APPLICATION NO. (If known, see 37 CFR 1.5) <b>09/744804</b>
INTERNATIONAL APPLICATION NO. <b>PCT/IL99/00417</b>	INTERNATIONAL FILING DATE <b>29 July 1999</b>	PRIORITY CLAIMED <b>30 July 1998</b>
TITLE OF INVENTION <b>TUMOR ASSOCIATEE ANTIGEN PEPTIDES AND USE OF SAME AS ANTI-TUMOR VACCINES</b>		
APPLICANT(S) FOR DO/EO/US <b>Lea EISENBACH et al.</b>		

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

- ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
- ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
- ☒ This is an express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).
- ☒ The US has been elected in a Demand by the expiration of 19 months from the priority date (PCT Article 31).
- ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
  - ☐ is attached hereto (required only if not transmitted by the International Bureau).
  - ☒ has been communicated by the International Bureau.
  - ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
- ☐ An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)).
- ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))
  - ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
  - ☐ have been communicated by the International Bureau.
  - ☐ have not been made; however, the time limit for making such amendments has NOT expired.
  - ☒ have not been made and will not be made.
- ☐ An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
- ☐ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
- ☐ An English language translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

**Items 11. to 16. below concern document(s) or information included:**

- ☒ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
- ☐ An Assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
- ☒ A **FIRST** preliminary amendment.
  - ☐ A **SECOND** or **SUBSEQUENT** preliminary amendment.
- ☐ A substitute specification.
- ☐ A change of power of attorney and/or address letter.
- ☒ Other items or information:
  - ☒ Courtesy copy of the International Application as filed.
  - ☒ Courtesy copy of the first page of the International Publication (WO 00/06723).
  - ☒ **Courtesy copy of the International Preliminary Examination Report with annexes containing claims 1-52 to be substituted for the original claims for examination in this case.**
  - ☒ Formal drawings, 29 sheets, Figures 1-24c.
  - ☒ Courtesy Copy of the International Search Report.

U.S. APPLICATION NO (If known, see 37 CFR 1.5) - <b>09/744804</b>		International Application No PCT/IL99/00417		Attorney's Docket No <b>EISENBACH 3</b>	
--	--	--	--	--	--

17. [xx] The following fees are submitted:

**BASIC NATIONAL FEE (37 CFR 1.492 (a)(1)-(5):**  
 Neither international preliminary examination fee (37 CFR 1.482)  
 nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO  
 and International Search Report not prepared by the EPO or JPO.....**\$1000.00**

International preliminary examination fee (37 CFR 1.482) not paid to  
 USPTO but International Search Report prepared by the EPO or JPO.....**\$860.00**

International preliminary examination fee (37 CFR 1.482) not paid to USPTO but  
 international search fee (37 CFR 1.445(a)(2)) paid to USPTO.....**\$710.00**

International preliminary examination fee paid to USPTO (37 CFR 1.482)  
 but all claims did not satisfy provisions of PCT Article 33(1)-(4)..... **\$690.00**

International preliminary examination fee paid to USPTO (37 CFR 1.482)  
 and all claims satisfied provisions of PCT Article 33(1)-(4).....**\$100.00**

**ENTER APPROPRIATE BASIC FEE AMOUNT =**

Surcharge of **\$130.00** for furnishing the oath or declaration later than [ ] 20 [X] 30  
 months from the earliest claimed priority date (37 CFR 1.492(e)).

Claims as Originally Presented	Number Filed	Number Extra	Rate		
Total Claims	49 - 20	29	X \$18.00	\$ 522.00	
Independent Claims	4 - 3	01	X \$80.00	\$ 80.00	
Multiple Dependent Claims (if applicable)			+\$270.00	\$	
<b>TOTAL OF ABOVE CALCULATIONS =</b>				<b>\$1,592.00</b>	

Claims After Post Filing Prel. Amend	Number Filed	Number Extra	Rate		
Total Claims	- 20		X \$18.00	\$	
Independent Claims	- 3		X \$78.00	\$	
<b>TOTAL OF ABOVE CALCULATIONS =</b>				<b>\$1,592.00</b>	

Reduction of ½ for filing by small entity, if applicable. Applicant claims small entity  
 status. See 37 CFR 1.27.

**SUBTOTAL =**

Processing fee of **\$130.00** for furnishing the English translation later than [ ] 20 [ ] 30  
 months from the earliest claimed priority date (37 CFR 1.492(f)).

**TOTAL NATIONAL FEE =**

Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be  
 accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31) **\$40.00** per property +

**TOTAL FEES ENCLOSED =**

	<b>Amount to be:</b>	\$
	<b>refunded</b>	
	<b>charged</b>	\$

**CALCULATIONS PTO USE ONLY**

a. [ ] A check in the amount of \$\_\_\_\_\_ to cover the above fees is enclosed.

b. [X] Credit Card Payment Form (PTO-2038), authorizing payment in the amount of \$ 796.00, is attached.


c. [ ] Please charge my Deposit Account No. **02-4035** in the amount of \$\_\_\_\_\_ to cover the above fees  
 A duplicate copy of this sheet is enclosed.

d. [XX] The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment  
 to Deposit Account No. **02-4035**. A duplicate copy of this sheet is enclosed.

**NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or  
 (b)) must be filed and granted to restore the application to pending status.**

SEND ALL CORRESPONDENCE TO:

**BROWDY AND NEIMARK, P.L.L.C.**  
**624 NINTH STREET, N.W., SUITE 300**  
**WASHINGTON, D.C. 20001**  
**TEL: (202) 628-5197**  
**FAX: (202) 737-3528**  
**Date of this submission: January 30, 2001**

  
 SIGNATURE  
Roger L. Browdy  
 NAME  
25,618  
 REGISTRATION NUMBER

09/744804

JCO7 Rec'd PCT/PTO 30 JAN 2001

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of:	)	Art Unit:
Lea EISENBACH et al.	)	
	)	
IA No.: PCT/IL99/00417	)	
	)	Washington, D.C.
IA Filed: 29 July 1999	)	
	)	
U.S. App. No.:	)	
(Not Yet Assigned)	)	
	)	January 30, 2001
National Filing Date:	)	
(Not Yet Received)	)	
	)	
For: TUMOR ASSOCIATED...	)	Docket No.:
		EISENBACH 3

PRELIMINARY AMENDMENT

Honorable Commissioner for Patents and Trademarks  
Washington, D.C. 20231

Sir:

Contemporaneous with the filing of this case and  
prior to calculation of the filing fee, kindly amend as  
follows:

IN THE SPECIFICATION

After the title please insert the following  
paragraph:

--The present application is the national stage  
under 35 U.S.C. 371 of PCT/IL99/00417, filed 29 July 1999.--

RECEIVED 10/11/00

IN THE CLAIMS

Claim 19, line 1, delete "any of claims 1-18", and insert therefor --claim 1--.

Claim 21, line 1, delete "any of claims 1-20", and insert therefor --claim 1--.

Claim 23, line 1, delete "or 22".

Claim 24, line 1, delete "any of claims 1-23", and insert therefor --claim 1--.

Claim 30, line 2, delete "any of claims 1-23", and insert therefor --claim 1--.

Claim 35, line 3, delete "any of claims 24-29", and insert therefor --claim 24--.

Claim 36, line 3, delete "any of claims 30-34", and insert therefor --claim 30--.

Claim 37, line 1, delete "any of claims 1-23", and insert therefor --claim 1--.

Claim 39, line 1, delete "or 38".

Claim 40, line 2, delete "any claims 37-39", and insert therefor --claim 38--.

Claim 41, line 2, delete "any of claims 1-23", and insert therefor --claim 1--.

Claim 45, line 1, delete "any of claims 24-29 or 40", and insert therefor --claim 24--.

Claim 47, line 1, delete "any of claims 30-34", and insert therefor --claim 30--.

Delete claims 49-51.

REMARKS

The above amendment to the specification is being made to insert reference to the PCT application of which the present case is a U.S. national stage. The above amendments to the claims are being made in order to eliminate any properly multiply dependent claims, for the purpose of reducing the filing fee. Please enter this amendment prior to calculation of the filing fee in this case.

Favorable consideration is earnestly solicited.

Respectfully submitted,  
BROWDY AND NEIMARK, P.L.L.C.  
Attorneys for Applicant

By: 

Roger L. Browdy  
Registration No. 25,618

RLB:wrđ  
Telephone No.: (202) 628-5197  
Facsimile No.: (202) 737-3528

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of:	)	Box PCT
EISENBACH et al	)	Examiner:
Appln. No.: 09/744,804	)	Washington, D.C.
Filed: January 30, 2001	)	June 20, 2001
For: TUMOR ASSOCIATED ANTIGEN	)	Atty.Docket: EISENBACH=3
.PEPTIDES AND USE OF SAME	)	
AS ANTI-TUMOR VACCINES	)	

RESPONSE TO NOTIFICATION TO COMPLY WITH  
SEQUENCE LISTING REQUIREMENTS

Honorable Commissioner for Patents  
Washington, D.C. 20231

Sir:

In response to the Notification to Comply dated  
April 30, 2001, and prior to the examination of the above-  
described application, please amend the present application as  
follows:

IN THE SPECIFICATION

Please replace the paragraph beginning at page 33,  
line 3, with the following rewritten paragraph:

--TABLE 3a  
*Predicted human Uroplakin Ib, II and III peptides that bind to HLA-A2*

Peptide	Start Position	Sequence	SEQ ID NO:
Uroplakin Ib/B1	239	AILCWTFWV	50
Uroplakin Ib/B2	92	FILMFIVYA	51
Uroplakin Ib/B3	29	LTAECIFFV	52
Uroplakin Ib/B4	154	MLQDNCCGV	53
Uroplakin Ib/B5	240	ILCWTFWVL	54
Uroplakin Ib/B6	86	KILLAYFIL	55
Uroplakin Ib/B7	64	FVGICLFCL	56
Uroplakin II/8	161	VLLSVAMFL	57
Uroplakin II/9	162	LLSVAMFLL	58
Uroplakin III/3.1	214	ILGSLPFFL	59
Uroplakin III/3.2	128	ILNAYLVRV	60
Uroplakin III/3.3	221	FLLVGFAGA	61
Uroplakin III/3.4	20	NLQPQLASV	62
Uroplakin III/3.5	47	CMFDSKEAL	63
Uroplakin III/3.6	62	YLYVLVDSA	64
Tyrosinase	368	YMDGTMSQV	65

Table 3a shows the sequences of the peptides tested in single letter amino acid code and their starting position in the intact protein (according to NCBI accession nos. 3298345 (peptides 3.1-3.6), 3483011 (peptides 8 and 9), and 3721858 (peptides B1-B7)).--

Please replace the paragraph beginning at page 43, line 21, with the following rewritten paragraph:

--TABLE 9  
*Predicted human Cripto-1 derived peptides that bind to HLA-A2*

Peptide	Start Position	Sequence	SEQ ID NO:
Cripto-1/C1	5	KMARFSYSV	66
Cripto-1/C2	151	GLVMDEHLV	67
Cripto-1/C3	145	FLPGCDGLV	68
Cripto-1/C4	89	CMLGSFCAC	69
Cripto-1/C5	43	YLAFRDDSI	70
Cripto-1/C6	123	WLPKKCSLC	71
Cripto-1/C7	83	CLNGGTCML	72
Cripto-1/C8	176	MLVGICLSI	73
Cripto-1/C9	23	FELGLVAGL	74
Cripto-1/C10	5	KMVRFSYSV	75
Cripto-1/C11	83	CLNEGTCML	76
Cripto-1/C12	176	MLAGICLSI	77

Table 9 shows the sequences of the peptides tested in single letter amino acid code and their starting position in the intact protein (according to NCBI accession nos. 117473 (C1-C9) and 321120 (C10-C12)).--



REMARKS

Responsive to the Notification to Comply with Sequence Listing Requirements, attached hereto is a 3 1/2" disk containing the "Sequence Listing" in computer readable form in accordance with 37 C.F.R. §1.821(e).

Applicants have amended the specification to insert SEQ ID Nos, as supported in the present specification.

The following statement is provided to meet the requirements of 37 C.F.R. §1.821(f) and 1.821(g).

I hereby state, in accordance with 37 C.F.R. §1.821(f), that the content of the paper copy sequence listing as filed and the attached computer readable copy of the sequence listing are believed to be the same.

I hereby also state, in accordance with 37 C.F.R. §1.821(g), that the submission is not believed to include new matter.

Under U.S. rules, each sequence must be classified in <213> as an "Artificial Sequence", a sequence of "Unknown" origin, or a sequence originating in a particular organism, identified by its scientific name.

Neither the rules nor the MPEP clarify the nature of the relationship which must exist between a listed sequence and an organism for that organism to be identified as the origin of the sequence under <213>.

Hence, counsel may choose to identify a listed sequence as associated with a particular organism even though that sequence does not occur in nature by itself in that

organism (it may be, e.g., an epitopic fragment of a naturally occurring protein, or a cDNA of a naturally occurring mRNA, or even a substitution mutant of a naturally occurring sequence). Hence, the identification of an organism in <213> should not be construed as an admission that the sequence *per se* occurs in nature in said organism.

Similarly, designation of a sequence as "artificial" should not be construed as a representation that the sequence has no association with any organism. For example, a primer or probe may be designated as "artificial" even though it is necessarily complementary to some target sequence, which may occur in nature. Or an "artificial" sequence may be a substitution mutant of a natural sequence, or a chimera of two or more natural sequences, or a cDNA (i.e., intron-free sequence) corresponding to an intron-containing gene, or otherwise a fragment of a natural sequence.

The Examiner should be able to judge the relationship of the enumerated sequences to natural sequences by giving full consideration to the specification, the art cited therein, any further art cited in an IDS, and the results of his or her sequence search against a database containing known natural sequences.

Attached hereto is a marked-up version of the changes made to the specification and claims by the current amendment. The attached page is captioned "Version with markings to show changes made".


Applicants submit that the present application contains patentable subject matter and therefore urge the examiner to pass the case to issuance.

If the examiner has any questions or comments concerning the above described application, the examiner is urged to contact the undersigned at the phone number below.

Respectfully submitted,

BROWDY AND NEIMARK, P.L.L.C.  
Attorneys for Applicant(s)

By

  
ROGER L. BROWDY  
Registration No. 25,618

RLB:al  
624 Ninth Street, N.W.  
Washington, D.C. 20001  
Telephone No.: (202) 628-5197  
Facsimile No.: (202) 737-3528

F:\,Y\YEDA\Eisenbach 3\PTO\RESPONSE TO NOTICE TO COMPLY.doc

VERSION WITH MARKINGS TO SHOW CHANGES MADE

In the specification:

The paragraph beginning at line 3 of page 33 has been amended as follows:

TABLE 3a  
*Predicted human Uroplakin Ib, II and III peptides that bind to HLA-A2*

Peptide	Start Position	Sequence	<u>SEQ ID NO.:</u>
Uroplakin Ib/B1	239	AILCWTFWV	<u>50</u>
Uroplakin Ib/B2	92	FILMFIVYA	<u>51</u>
Uroplakin Ib/B3	29	LTAECIFFV	<u>52</u>
Uroplakin Ib/B4	154	MLQDNCCGV	<u>53</u>
Uroplakin Ib/B5	240	ILCWTFWVL	<u>54</u>
Uroplakin Ib/B6	86	KILLAYFIL	<u>55</u>
Uroplakin Ib/B7	64	FVGICLFCL	<u>56</u>
Uroplakin II/8	161	VLLSVAMFL	<u>57</u>
Uroplakin II/9	162	LLSVAMFLL	<u>58</u>
Uroplakin III/3.1	214	ILGSLPFFL	<u>59</u>
Uroplakin III/3.2	128	ILNAYLVRV	<u>60</u>
Uroplakin III/3.3	221	FLLVGFAGA	<u>61</u>
Uroplakin III/3.4	20	NLQPQLASV	<u>62</u>
Uroplakin III/3.5	47	CMFDSKEAL	<u>63</u>
Uroplakin III/3.6	62	YLYVLVDSA	<u>64</u>
Tyrosinase	368	YMDGTMSQV	<u>65</u>

Table 3a shows the sequences of the peptides tested in single letter amino acid code and their starting position in the intact protein (according to NCBI accession nos. 3298345 (peptides 3.1-3.6), 3483011 (peptides 8 and 9), and 3721858 (peptides B1-B7)).

The paragraph beginning at line 21 of page 43 has been amended as follows:

TABLE 9  
Predicted human Cripto-1 derived peptides that bind to HLA-A2

Peptide	Start Position	Sequence	SEQ ID NO.
Cripto-1/C1	5	KMARFSYSV	66
Cripto-1/C2	151	GLVMDEHLV	67
Cripto-1/C3	145	FLPGCDGLV	68
Cripto-1/C4	89	CMLGSFCAC	69
Cripto-1/C5	43	YLAFRDDSI	70
Cripto-1/C6	123	WLPKKCSLC	71
Cripto-1/C7	83	CLNGGTCML	72
Cripto-1/C8	176	MLVGICLSI	73
Cripto-1/C9	23	FELGLVAGL	74
Cripto-1/C10	5	KMVRFSYSV	75
Cripto-1/C11	83	CLNEGTCML	76
Cripto-1/C12	176	MLAGICLSI	77

Table 9 shows the sequences of the peptides tested in single letter amino acid code and their starting position in the intact protein (according to NCBI accession nos. 117473 (C1-C9) and 321120 (C10-C12)).

JC07 Rec'd PCT/PTO 30 JAN 2001

TUMOR ASSOCIATED ANTIGEN PEPTIDES AND USE OF SAME AS  
ANTI-TUMOR VACCINES

FIELD AND BACKGROUND OF THE INVENTION

5 The present invention relates to tumor associated antigen (TAA) peptides and to the use of same, of polynucleotides encoding same and of cells presenting same as anti-tumor vaccines. More particularly, the present invention relates to tumor associated antigen peptides derived from Uroplakin Ia, Ib, II and III, Prostate specific antigen (PSA), Prostate acid phosphatase (PAP) and Prostate  
10 specific membrane antigen (PSMA), BA-46 (Lactadherin), Mucin (MUC-1), and Teratocarcinoma-derived growth factor (CRIPTO-1) and the use of same as anti-tumor vaccines to prevent or cure bladder, prostate, breast or other cancers, carcinomas in particular. Most particularly, the present invention relates to tumor associated antigen peptides which are presentable to the immune system by  
15 HLA-A2-molecules.

Local therapy such as surgical excision or ablation by radiation is a mainstay for the treatment of primary cancer and is curative for a percentage of patients. However, many malignancies will recur locally or at a distant site. Thus the prevention or cure of metastases remains a major focus in clinical oncology  
20 (1). Although early detection followed by surgery provides good prognosis for a number of major cancer types, a large fraction of patients would need adjuvant therapy. Part of these patients will, with time, succumb to metastasis (2-4). Alternative approaches based on gene therapy and immunotherapy have been the focus of attention in the last years. One such approach is specific active  
25 immunotherapy (SAI, 5). The objective of SAI is to stimulate a tumor specific cytotoxic T lymphocytes (CTL) immune response that is capable of eliminating residual metastatic disease and induce a state of immunity to protect the patients from recurrent disease. The underlying assumption of SAI is that tumor cells express tumor antigens which are sufficiently distinct in structure or context to  
30 induce an effective CTL response (6). Although the validity of these assumptions was questioned, a number of studies in the last decade have demonstrated the rational of SAI. In a landmark study, van Pel and Boon have shown that tumor associated antigens (TAAs) can be isolated and defined (7). Importantly, *ex-vivo* manipulations of "non-immunogenic" animal tumor cells can be used to elicit  
35 effective immune responses which will also recognize parental "non-immunogenic" tumor cells (8). Studies employing rodent tumor models with little intrinsic immunogenicity have shown that genetically modified tumor cells transduced to express MHC class I, cytokines such as IL-1, IL-2, IL-4, IL-6, IL-7, IL-12, IFN or GM-CSF or costimulatory molecules such as B7-1 or B7-2 were

capable of preventing or causing regression of tumors or metastases (reviewed in 9). Although gene modified tumor vaccine (GMTV) clinical trials, with improved retroviral vectors or other transfer methodologies are currently tested, it becomes clear that GMTV using autologous tumor cells might be limited by its complexity, high cost and ineffective gene transfer methodologies (10). One alternative approach would be vaccination with tumor associated antigens (TAAs) presented in an effective way to the patient's immune system, to induce antigen specific CTL (11).

Cytotoxic T lymphocytes (CTL), directed against peptides presented by MHC class I molecules, constitute powerful effectors of the immune system against tumors or infectious agents (12). These peptides are usually 8-10 amino acids long with 2-3 primary anchor residues that interact with the MHC class I molecules and 2-3 amino acid residues which bind to the T cell receptor (13). Several methods have been employed to identify CTL epitopes. If the amino acid sequence of a protein antigen is known, like in the case of viral proteins, oncogenes, suppressor genes or growth factor receptors, overlapping peptides of 8-10 amino acids in length can be synthesized and screened as CTL targets (14). CTL epitopes may also be identified subsequent to the search for MHC binding motifs in known proteins (15). If the tumor antigen is not known, isolation of the TAA peptides from total acid extract or from MHC class I molecules followed by HPLC fractionation steps and Edman sequencing (16) or mass spectrometry (17) provide a direct way of identifying CTL epitopes. Recently, a synthetic combinatorial library approach, in which defined amino acids in two MHC anchor positions are fixed and all other positions are subgrouped for CTL screening has led to the description of novel EL4 TAA peptide mimotopes (18).

The most fruitful method, so far, designed by T. Boon and his colleagues is the genetic approach in which cDNA expression libraries are pool transfected into COS7 cells with the appropriate HLA and screened by CTL lines. This approach led to the discovery of several human melanoma and mouse mastocytoma antigens recognized by specific CTL (19). The first report of a phase I clinical trial with the synthetic MAGE3 melanoma peptide, restricted by HLA-A1, showed regression of cutaneous, subcutaneous and lung metastases in 3/6 patients (20). Recently, two reports of clinical trials have shown that treatment of patients with a melanoma gp100 TAA peptide together with IL-2 resulted in significant tumor regression in 13/31 (42 %) patients and that vaccination with defined peptides or total peptide extracts on autologous dendritic cells (DC) resulted in complete or partial cures (21, 22). Regression of lung carcinoma established metastases or small established tumors was demonstrated in a murine model by peptide

vaccination (23, 24). These observations suggest that TAA peptide vaccines may constitute a reasonable therapeutic modality in advanced cancer. In studies with murine tumors, CTL are induced *in vivo* by immunization with irradiated tumor cells, often gene modified by MHC class I; cytokine or costimulatory molecules like B7-1 or B7-2 genes (16, 18, 25). In melanomas, CTL lines were mostly induced from peripheral blood mononuclear cells (PBMC) of patients or from tumor infiltrated lymphocytes (TIL, 19, 26). Yet, most metastatic tumors are non-immunogenic tumors and it is extremely difficult to derive CTL lines or clones from TIL or patient's PBL. Moreover, *in vitro* propagated CTL clones do not always represent dominant anti-tumor specificities but rather sporadic clones surviving culture conditions. Lately, a number of studies have compared the CTL repertoire of viral or other defined peptides, restricted by HLA-A2.1 in human PBL from HLA-A2.1 expressing patients to CTL induced in HLA-A2.1 transgenic mice. Good concordance between human HLA-A2.1 and murine transgenic HLA-A2.1 CTL repertoire was found, confirming the potential of such transgenics in identification of human CTL epitopes (27). Although vaccination with defined peptides of HLA transgenic mice shows an overlapping repertoire to human CTL, vaccination of such mice with multi-epitope proteins shows that murine H-2 restricted responses are dominant and obliterate, as a rule, cytolytic responses with direct recognition of human HLA (28). Thus, by combining classical HLA class I transgenesis with selective destruction of murine H-2, it is possible to derive useful mouse strains for the study of HLA class I restricted responses. While reducing the present invention to practice, we utilized such mice for preparation of anti-tumor CTL as a tool for TAA purification and as a model system to assess the immunogenicity of peptides.

Murine H-2 knockout mice transgenic for a single human HLA seem to be a suitable model for induction of anti-tumor CTL. Classical  $\beta_2$  microglobulin knockout mice ( $\beta_2m^{-/-}$ ) do not express H-2K<sup>b</sup> or other non-classical class I molecules, yet they express low levels of H-2D<sup>b</sup> heavy chain in the absence of  $\beta_2m$ . To derive fully H-2 knockout mice, Prof. F. Lemonnier (Pasteur Institute, Paris), prepared H-2D<sup>b</sup> $^{-/-}$  mice. These mice were crossed with  $\beta_2m^{-/-}$  mice and bred to derive homozygous  $\beta_2m^{-/-}$ , D<sup>b</sup> $^{-/-}$  mice that do not express any H-2 class I. These mice are practically depleted of CD8<sup>+</sup> splenocytes, as well as other CD8<sup>+</sup> cells. To reconstitute in these mice expression of a stable HLA-A2.1, expression of  $\beta_2m$  is necessary. A construct containing a leader sequence, domains  $\alpha_1$  and  $\alpha_2$  of HLA-A2.1 and  $\alpha_3$ , transmembrane and cytoplasmic domains of H-2D<sup>b</sup> fused to human  $\beta_2m$  (HhD) was prepared. The exchange of the  $\alpha_3$  human domain by a murine domain in HhD is thought to improve the interaction of the class I



molecule with CD8 molecules of the murine CTL (29). This HhD construct was transfected into RMA and RMA-S cells and shown to bind HLA-A2.1 restricted peptides. The HhD construct was used to produce transgenic mice in C57BL/6 recipients and positive founder mice were bred to the  $\beta 2m^{-/-}$ ,  $D^b^{-/-}$  mice (30).

5       The  $\beta 2m^{-/-}$ ,  $D^b^{-/-}$ , HhD $^{+/+}$  heterozygous mice show reconstitution of CD8 $^{+}$  cells in the periphery relative to  $\beta 2m^{-/-}$   $D^b^{-/-}$  mice. Moreover, preliminary data from Prof. Lemonnier's lab showed that CTL induced in HhD mice against influenza NP are directed to the same HLA-A2 dominant epitope as in the human repertoire. Homozygous HhD mice were derived and a colony was established in  
10       the Weizmann Institute of Science, Israel.

#### SUMMARY OF THE INVENTION

According to one aspect of the present invention there is provided a peptide derived from a protein selected from the group consisting of Uroplakin (UP),  
15       Prostate specific antigen (PSA), Prostate specific membrane antigen (PSMA), Prostate acid phosphatase (PAP), Lactadherin (BA-46) and Mucin (MUC1), and Teratocarcinoma-derived growth factor (CRIPTO-1), the peptide comprising 8 to 10 amino acid residues, of which a second residue from an amino terminal of the peptide and a carboxy terminal residue of the peptide are hydrophobic or  
20       hydrophilic natural or non natural amino acid residues, for example, SEQ ID NOs:1-64 and 65 to 77.

According to further features in preferred embodiments of the invention described below, the peptide is derived from Uroplakin, such as Uroplakin II, Uroplakin Ia, Uroplakin III and Uroplakin Ib, for example, SEQ ID NOs:1-19 or  
25       50-64.

According to still further features in the described preferred embodiments the peptide is derived from the Prostate specific antigen (PSA), for example, SEQ ID NOs:20-24.

According to still further features in the described preferred embodiments the peptide is derived from the Prostate specific membrane antigen (PSMA), for  
30       example, SEQ ID NOs:25-30.

According to still further features in the described preferred embodiments the peptide is derived from the Prostate acid phosphatase (PAP), for example, SEQ ID NOs:31-34.

35       According to still further features in the described preferred embodiments the peptide is derived from the Mucin, for example, SEQ ID NOs:42-49.

According to still further features in the described preferred embodiments the peptide is derived from a non tandem repeat array of the Mucin.

According to still further features in the described preferred embodiments the peptide is derived from a region selected from the group consisting of a signal peptide, a cytoplasmic domain and an extracellular domain of the Mucin.

According to still further features in the described preferred embodiments the peptide is derived from the Lactadherin (BA-46), for example, SEQ ID NOs:35-41.

According to still further features in the described preferred embodiments, the peptide is derived from the Teratocarcinoma-derived growth factor (CRIPTO-1), for example, SEQ ID NOs. 66 to 77.

According to still further features in the described preferred embodiments the Uroplakin (UP), Prostate specific antigen (PSA), Prostate specific membrane antigen (PSMA), Prostate acid phosphatase (PAP), Lactadherin (BA-46) and Mucin (MUC1) and Teratocarcinom-derived growth factor (CRIPTO-1) are each independently of mammalian origin.

According to still further features in the described preferred embodiments the mammal is selected from the group consisting of a humanoid and a rodent.

According to still further features in the described preferred embodiments the peptide includes at least one non-natural modification.

According to still further features in the described preferred embodiments the peptide includes at least one non-natural modification rendering peptides more immunogenic or more stable.

According to still further features in the described preferred embodiments the at least one modification is selected from the group consisting of peptoid modification, semipeptoid modification, cyclic peptide modification, N terminus modification, C terminus modification, peptide bond modification, backbone modification and residue modification.

According to another aspect of the present invention there is provided a pharmaceutical composition comprising, as an active ingredient, at least one peptide at set forth herein and a pharmaceutically acceptable carrier.

According to another aspect of the present invention there is provided a vaccine composition comprising, as an active ingredient, at least one peptide at set forth herein and a vaccination acceptable carrier.

According to still further features in the described preferred embodiments the carrier is selected from the group consisting of a proteinaceous carrier to which the at least one tumor associated antigen peptide is linked, an adjuvant, a protein or a recombinant protein and an antigen presenting cell.

According to still further features in the described preferred embodiments the composition, pharmaceutical or vaccine, is effective in prevention or cure of cancer or carcinoma metastases.

According to still further features in the described preferred embodiments the cancer is selected from the group consisting of breast, bladder, prostate, pancreas, ovary, thyroid, colon, stomach and head and neck cancer.

According to still further features in the described preferred embodiments the cancer is a carcinoma.

Further according to the present invention there is provided a method of prevention or cure of a cancer or of metastases thereof comprising the step of administering to a patient an effective amount of the pharmaceutical composition described herein.

Still further according to the present invention there is provided a method of prevention or cure of a cancer or of metastases thereof comprising the step of vaccinating a patient with an effective amount of the vaccine composition described herein.

According to yet another aspect of the present invention there is provided a polynucleotide encoding at least one peptide as set forth herein. One ordinarily skilled in the art would know how to reverse translate any of the peptides according to the present invention such as the peptides set forth in SEQ ID NOs:1-64 and 66 to 77, using, for example, the human preferred codon usage, to derive the sequences of the polynucleotides according to the present invention. Such polynucleotides are readily synthesizable using the well known solid phase technology for preparation of oligonucleotides such as oligodeoxynucleotides or analogs thereof.

According to still further features in the described preferred embodiments the polynucleotide forms a part of a longer polynucleotide designed to encode a fused protein product from which at least one peptide is cleavable by a protease.

According to a further aspect of the present invention there is provided a pharmaceutical composition comprising, as an active ingredient, at least one polynucleotide as set forth herein and a pharmaceutically acceptable carrier.

According to still a further aspect of the present invention there is provided a cellular vaccine composition comprising an antigen presenting cell presenting at least one peptide derived from a protein selected from the group consisting of Uroplakin (UP), Prostate specific antigen (PSA), Prostate specific membrane antigen (PSMA), Prostate acid phosphatase (PAP), Lactadherin (BA-46) Mucin (MUC1) and Teratocarcinoma-derived growth factor (CRIPTO-1), the at least one peptide comprising 9 or 10 amino acid residues, of which a second residue from

an amino terminal of the peptide and a carboxy terminal residue of the peptide are hydrophilic or aliphatic non-charged natural or synthetic amino acid residues.

According to still further features in the described preferred embodiments the antigen presenting cell is selected from the group consisting of a dendritic cell,  
5 a macrophage, a B cell and a fibroblast.

According to still further features in the described preferred embodiments the antigen presenting cell is caused to present the at least one tumor associated antigen peptide by a method selected from the group consisting of (a) genetically  
10 modifying the antigen presenting cell with at least one polynucleotide encoding the at least one tumor associated antigen peptide such that the said peptide or at least one longer polypeptide including said peptide will be expressed; (b) loading the antigen presenting cell with at least one polynucleotide encoding the at least one tumor associated antigen peptide; (c) loading the antigen presenting cell with the at least one tumor associated antigen peptide; and (d) loading the antigen  
15 presenting cell with at least one longer polypeptide including the at least one tumor associated antigen peptide.

According to a still further aspect of the present invention there is provided a pharmaceutical composition also comprising a helper peptide.

According to a still further aspect of the present invention there is provided  
20 a pharmaceutical composition, wherein the helper peptide has a T helper epitope.

According to a still further aspect of the present invention there is provided a vaccine composition also comprising a helper peptide.

According to a still further aspect of the present invention there is provided a vaccine composition, wherein the helper peptide has a T helper epitope.

25 According to a still further aspect of the present invention there is provided a use of the at least one peptide in the manufacture of a medicament.

According to a still further aspect of the present invention there is provided the at least one peptide for use as a medicament.

30 According to a still further aspect of the present invention there is provided a use of the at least one peptide in the manufacture of a medicament for the prevention or cure of a cancer or cancer metastases.

According to a still further aspect of the present invention there is provided the at least one peptide for use as a medicament for the prevention or cure of a cancer or cancer metastases.

35 According to a still further aspect of the present invention there is provided a peptide derived from a protein selected from the group consisting of Uroplakin (UP), Prostate specific antigen (PSA), Prostate specific membrane antigen (PSMA), Prostate acid phosphatase (PAP), Lactadherin (BA-46) and Mucin

(MUC1) and Teratocarcinoma-derived growth factor (CRIPTO-1), the peptide comprising 8-10 amino acid residues as selected so as to promote effective binding to a MHC class 1 type molecule such that a CTL response is elicitable.

The present invention successfully addresses the shortcomings of the presently known configurations by providing novel tumor associated antigen peptides effective in eliciting CTL response which may therefore be effective therapeutic agents to combat cancer.

#### BRIEF DESCRIPTION OF THE DRAWINGS

10 The invention herein described, by way of example only, with reference to the accompanying drawings, wherein:

FIG. 1 demonstrates stabilization of cell surface HhD on RMA-S cells by human Uroplakin II derived peptides. Uroplakin II derived synthetic peptides (HURO1-7) or a peptide from the melanoma associated TAA Tyrosinase (Tyr) which served as a control were loaded at various concentrations (3-300  $\mu$ M) on the RMA-HhD cells as described. Indirect fluorescence activated cell sorter (FACS) analyses were performed by incubating  $5 \times 10^5$  loaded cells with the anti-HLA-A2 monoclonal antibodies BB7.2 for 30 minutes at 4° C. After washing the cells in PBS-0.5 %, BSA-0.1 %, sodium azide, the second antibody, goat anti mouse-FITC was applied for 30 minutes at 4° C. Following another wash the fluorescence was recorded on a FACS scan (BD). Mean fluorescence at 300  $\mu$ M is shown.

FIG. 2 demonstrates lytic activity of human peripheral blood lymphocytes (PBL) against Uroplakin II loaded target cells. PBL from HLA-A2 positive patients were resensitized with tumor extracted peptide preparations from transitional cell carcinoma (TCC) specimens as described in Materials and Methods. Resensitized lymphocytes from 4 TCC patients treated with Bacillus Calmette Guerein (BCG), 2 non-treated TCC patients, and 1 prostate cancer patient (control), were admixed at different ratios with labeled T2 cells, loaded with TCC extracted peptides (TCC pool) normal bladder mucosa extracted peptides (bladder), synthetic Uroplakin II peptides (HURO1-7) or the melanoma peptide from Tyrosinase (Tyr). Labeled target cells, J82, a HLA-A2 expressing TCC line, SUP, a low class I expressor and K562, a NK sensitive line were also tested. Percent specific lysis, at effector: target of 100:1 is presented.

FIG. 3 demonstrates lytic activity of human peripheral blood lymphocytes against peptides extracted from individual TCC tumors. PBL were resensitized as in Figure 2. Peptides extracted from 9 individual tumors (TCC1-9), that were

pooled for resensitization, were loaded on T2 cells. Labeled loaded targets were tested for lysis by sensitized PBL as described in Figure 2.

FIG. 4 demonstrates lysis of human Uroplakin II and murine Uroplakin II peptide loaded target cells by CTL induced against human Uroplakin II derived peptides in HhD mice. Mice were immunized as described, by RMA-S-HhD-B7.1 cells, loaded with human Uroplakin II derived synthetic peptides 1-7 as described in Materials and Methods. Loading was performed with individual peptides and cells were pooled for immunization. Resensitized CTL (see Materials and Methods) were tested for lysis of RMA-S-HhD targets loaded with human or murine Uroplakin II peptides (HURO 1-7, MURO1, 3, 4, 6) with TCC or normal bladder extracted peptides (RMA-S TCC, RMA-S bladder), with the tyrosinase peptide (RMA-S Tyr) with peptides derived from breast tumors (RMA-S Br.) or with the cell lines J82 or SUP. E:T ratios of 100:1, 50:1, 25:1 and 12.5:1 were tested and the results are presented in lytic units LU30 was calculated by linear regression analysis of percentage lysis vs. the log (ln) of effector cell number and expressed as the number of LU per  $10^6$  effector cells. The coefficient was 0.85 - 0.95 for all groups.

FIG. 5 demonstrates lysis of human Uroplakin II and Ia and murine Uroplakin II derived peptide loaded target cells by CTL induced against TCC extracted peptides in HhD mice. HhD mice were immunized by RMA-S-HhD-B7.1 cells loaded with peptides extracted from 9 TCC samples and pooled. CTL were prepared as described and tested for lysis of labeled RMA-S-HhD target cells loaded with synthetic, tissue derived or control peptides and cell lines as in Figure 4 and synthetic peptides derived from Uroplakin Ia (HURO 11-18). The results are presented in lytic units as described in Figure 4.

FIG. 6 demonstrates stabilization of cell surface HhD on RMA-S cells by human Prostate specific antigen (PSA) and Prostate acid phosphatase (PAP) derived peptides. PSA and PAP derived peptides were loaded at various concentrations (1  $\mu$ M - 1 mM) on RMA-S-HhD cells. FACS analysis was performed as described in Figure 1, wherein Tyr served as a control. Mean fluorescence at 1 mM is presented.

FIG. 7 demonstrates stabilization of cell surface HhD on RMA-S cells by human Prostate specific membrane antigen (PSMA) derived peptides. PSMA derived peptides (1  $\mu$ M - 1 mM) were loaded and cells were monitored as described in Figure 6.

FIG. 8 demonstrates immunogenicity and antigenicity of PAP derived peptides in HhD mice. HhD mice were immunized as described, by RMA-S-HhD-B7.1 cells loaded individually with each PAP peptide and pooled

before vaccination. CTL were tested against RMA-HhD cells loaded with synthetic peptides (PAP 1-4), prostate carcinoma extracted peptides (RMA-S, CAP), hyperplastic (normal) prostate extracted peptides (RMA-S HP), breast carcinoma derived peptides (RMA-S Br, control) and tyrosinase (RMA-S Tyr, control). The prostate carcinoma cell line DU145 (DU) and its HhD transfectant (DU HhD) were also tested as targets. The results are presented in lytic units as in Figure 4.

FIG. 9 demonstrates immunogenicity and antigenicity of PSA derived peptides in HhD mice. Procedures were performed and results are presented as in Figure 8.

FIG. 10 demonstrates immunogenicity and antigenicity of PSMA derived peptides in HhD mice. Procedures were performed and results are presented as in Figure 8.

FIG. 11 demonstrates lysis of PSA, PAP and PSMA derived peptide loaded RMA-S-HhD cells by CTL induced against DU145-HhD cells in HhD mice. Mice were immunized with irradiated DU145-HhD cells and CTL assays were performed as described in Materials and Methods. Labeled RMA-S-HhD target cells were loaded with synthetic PSA (1-5), PAP (1-4), PSM-A (1-6) or tyrosinase (RMA-S Tyr) peptides, with prostate carcinoma (RMA-S CAP), breast carcinoma (RMA-S Br.) or hyperplastic prostate (RMA-S HP) extracted peptides. DU145 (DU) and DU145-HhD also served as targets. The results are presented in lytic units as in Figure 4.

FIG. 12 demonstrates lysis of PSA, PAP and PSMA derived peptide loaded RMA-S-HhD by CTL induced against prostate carcinoma extracted peptides in HhD mice. HhD mice were immunized with RMA-S-HhD-B7.1 cells loaded with peptides extracted from prostate carcinoma samples. Experimental details are equivalent to the details in Figure 11.

FIG. 13 demonstrates stabilization of cell surface HhD on RMA-S- cells by human BA-46 (Lactadherin) derived peptides. RMA-S-HhD cells were incubated with 1-100  $\mu$ M synthetic peptides and monitored as described in Figure 1.

FIG. 14 demonstrates immunogenicity of BA-46 peptides in HhD mice. Mice were immunized as described in Materials and Methods, with RMA-S-HhD-B7.1 cells loaded with synthetic peptides. CTL assays were performed on RMA-S-HhD target cells loaded with homologous peptides.

FIG. 15 demonstrates lysis of BA-46 peptide loaded targets by CTL induced against breast carcinoma extracted peptides in HhD mice. HhD mice were immunized with RMA-S-HhD-B7.1 cells loaded with peptides extracted

from breast carcinoma samples. Experimental details are as in Figure 11. The data is presented as percent specific lysis at E:T of 100:1.

FIG. 16 demonstrates differential lysis of tumor extracted peptide loaded target cells by CTL induced against tumor peptides or BA-46 derived peptides in HhD mice. Mice were immunized by RMA-S-HhD-B7.1 cells loaded with synthetic BA-46 derived peptides or breast carcinoma derived peptides as described in Materials and Methods. CTL were tested against RMA-S-HhD cells loaded with breast carcinoma extracted peptides or normal breast tissue extracted peptides. Percent specific lysis at E:T of 100:1 is presented.

FIG. 17 demonstrates HLA-A2.1 (HhD) restricted lysis of a breast carcinoma cell line by CTL induced against BA-46 derived peptides or tumor extracted peptides. HhD mice were immunized as in Figure 16 and CTL were tested against the HLA negative MDA-MB-157 and their HhD transfectants. The data is presented in percent specific lysis at a E:T of 100:1.

FIG. 18 demonstrates stabilization of cell surface HhD on RMA-S- cells by Mucin (MUC-1/A7, E6, D6) derived peptides. Details are as described in Figure 1.

FIG. 19 demonstrates immunogenicity of MUC-1 derived peptides. HhD mice were immunized and CTL tested as in Figure 14.

FIG. 20 demonstrates HLA-A2.1 (HhD) restricted lysis of a breast carcinoma cell line by CTL induced against MUC-1 derived peptides. Details as described in Figure 17.

FIG. 21 demonstrates lysis of MUC-1 loaded target cells by CTL induced against breast carcinoma extracted peptides in HhD mice. Details are as described in Figure 15. E:T is 50:1.

FIG. 22 demonstrates differential lysis of tumor extracted peptide loaded target cells by CTL induced against MUC-1 derived peptides in HhD mice. Details are as described in Figure 16.

FIGs. 23a-e demonstrate lysis of additional human Uroplakin (Ib, II and III) peptide loaded target cells by CTL induced against the peptides in HhD mice. Mice were immunized with RMA-S-HhD-B7.1 cells, loaded with Uroplakin derived synthetic peptides, as described in Materials and Methods. Loading was performed with individual peptides and cells were pooled for immunization. Resensitized CTL (see Materials and Methods) were tested for lysis of RMA-S-HhD targets loaded with Uroplakin peptides or with an irrelevant HLA-A2 binding peptide derived from tyrosinase. Results are expressed as % Lysis using E:T ratios of 100:1, 50:1, 25:1, and 12.5:1. In Figs. 23a-d peptides were loaded onto target cells at a concentration of 200  $\mu$ M. For Fig. 23e two



concentrations of peptide were used for loading (20  $\mu$ M and 200  $\mu$ M). The protein origins of the peptides examined are listed in Table 3.

FIG. 24a-c demonstrate lysis of human CRIPTO-a peptide loaded target cells by CTL induced against peptides in HhD mice. Mice were immunized with RMA-S-HhD-B7.1 cells, loaded with CRIPTO-1 derived synthetic peptides, as described in Materials and Methods. Loading was performed with individual peptides and cells were pooled for immunization (pool of C1, C3, C5, and C7 for Fig. 24, pool of C2, C4, C6, and C8 for Fig. 24b, and pool of C10-12 for Fig. 24c). Resensitized CTL were tested for lysis of RMA-S-HhD targets loaded with CRIPTO-1 peptides at two concentrations (200 or 20  $\mu$ M), or with an irrelevant HLA-A2 binding peptide (C2 for Fig. 24a, C10 for Fig. 24b). Results are expressed a % Lysis using E:T ratios of 100:1, 50:1, 25:1, and 12.5:1.

## 15 DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present invention concerns peptides and pharmaceutical and vaccine compositions including same which can be used to prevent or cure cancer, both primary tumors and metastases. Specifically, the present invention can be used to provide vaccines which include novel and potent tumor associated antigen peptides derived from Uroplakins, Prostate specific antigen, Prostate acid phosphatase, Prostate specific membrane antigen, Lactadherin Mucin and Teratocarcinoma-derived growth factor, which can be used as anti-tumor vaccines to prevent or cure, for example, bladder, prostate or breast cancers, carcinomas in particular, or any other tumor expressing the above listed proteins.

25 The principles and operation of the present invention may be better understood with reference to the drawings and accompanying descriptions.

Before explaining at least one embodiment of the invention in detail, it is to be understood that the invention is not limited in its application to the details of construction and the arrangement of the components set forth in the following description or illustrated in the drawings. The invention is capable of other 30 embodiments or of being practiced or carried out in various ways. Also, it is to be understood that the phraseology and terminology employed herein is for the purpose of description and should not be regarded as limiting.

A specific CTL response was found in patients with bladder transitional cell carcinoma (TCC) against the tumor extract and against peptides homologous to Uroplakin II sequence, which was significantly augmented after BCG intravesical therapy. All Uroplakin II peptides employed were good immunizers in HhD mice and CTL from these mice induced intense lysis of targets loaded

with Uroplakin II homologue peptides, TCC peptide extract and of the J82 HLA-A2 expressing cell line. CTL from HhD mice immunized with TCC peptide extract showed cross reactivity and induced significant lysis of targets loaded with Uroplakin II, or Uroplakin Ia homologue peptides, with TCC peptide extract and of J82 cells. For all immunized HhD mice, there was reduced lysis of targets loaded with normal bladder mucosa peptide extract and of TCC SUP cells (minimal HLA-A2 expression). Also there was no non-specific lysis. In addition, a cross reactivity of CTL induced by human Uroplakin II peptides against murine homologue peptides was found. Despite this finding there was no damage, or inflammatory infiltrate in the internal organs, including the urinary bladder on careful histological examination. These data indicate that Uroplakin homologue peptides constitute specific CTL epitopes and may be considered for immunotherapeutic vaccines alone, or in combination with intravesical BCG instillations in patients prone to recurrence and progression of bladder TCC. Similar results were obtained with human Uroplakin Ib and III.

By using HhD mice, HLA-A2 transgenic and which do not express murine MHC class I, it was found that 3 Prostate specific antigen (PSA), 4 Prostate specific membrane antigen (PSMA) and 4 Prostate acid phosphatase (PAP) homologue peptides were immunogenic. The 3 PSA homologue peptides and 2 of the 4 PSMA homologue peptides were not evaluated previously. The 2 PSMA homologue peptides, which were evaluated previously and were found to be CTL epitopes in CAP patients, were also found to be immunogenic in HhD mice. PAP homologue peptides were not evaluated previously and all 4 HLA-A2 binding peptides were very immunogenic in HhD mice and induced also, specific lysis of DU145-HhD cells. Moreover, CTL derived from HhD mice immunized with CAP peptide extract, or with DU145-HhD cells induced specific lysis of targets loaded with CAP peptide extract and PAP homologue peptides. There was also cross reactivity between CTL derived from CAP immunized HhD mice and PSA, or PSMA homologue peptide loaded targets. Low cross reactivity was found for CTL derived from DU145-HhD immunized HhD mice, probably due to low expression of these proteins by DU145 cells. There was no damage, or inflammatory infiltrate in the internal organs of immunized or control HhD mice on careful histological examination. These data indicate that PSA, PSMA and especially PAP homologue peptides constitute specific CTL epitopes and may be considered for immunotherapy of CAP patients.

It was further found a specific CTL response in HhD mice against breast tumor extract and against novel peptides complementary to Lactadherin (BA-46) sequences. These data indicate that BA-46 peptides constitute specific CTL

epitopes enriched in breast carcinoma and may be considered for immunotherapeutic vaccines.

It was yet further found that Mucin (MUC-1) derived peptides are potential TAA peptides that can be used in anti-tumor vaccine preparations. CTL induced by these peptides lyse better tumor extract loaded targets than normal tissue extract loaded targets. The lysis is HLA-A2 restricted and breast specific. Three novel peptides, from non-tandem repeat array (TRA) domains were shown to constitute potential CTL epitopes.

Thus, in accordance with one aspect of the teachings of the present invention there is provided a vaccine composition which includes at least one tumor associated antigen peptide derived from a protein selected from the group consisting of Uroplakin (UP), Prostate specific antigen (PSA), Prostate specific membrane antigen (PSMA), Prostate acid phosphatase (PAP), Lactadherin (BA-46) and Mucin (MUC1).

A specific CTL response has been obtained from Teratocarcinoma-derived growth factor (CRIPTO-1) derived peptide in HhD transgenic mice. This data indicates CRIPTO-1 peptides may constitute effective CTL epitopes in humans with HLA-A2 haplotypes and may be considered for immunotherapeutic cancer vaccines.

According to another aspect of the present invention there is provided a method of vaccination for prevention or cure of cancer. The method is effected by administering to a patient a vaccine composition including at least one tumor associated antigen peptide derived from a protein selected from the group consisting of Uroplakin (UP), Prostate specific antigen (PSA), Prostate specific membrane antigen (PSMA), Prostate acid phosphatase (PAP), Lactadherin (BA-46) and Mucin (MUC1).

Immunotherapy by *in vivo* DNA transfer of DNA coding for TAA is based on the rationale of quality or quantity increased peptide presentation leading to activation of an immune response against these peptides. Gene or DNA vaccination results in the intracellular processing and presentation of immunogenic peptides (99). Initial reports on DNA vaccination showed that "naked" DNA injected into the muscle tissue of a mouse is expressed efficiently (32). Embryonically expressed TAA such as CEA was tested (33). Immunization of mice with CEA expressing plasmid DNA was indeed found to protect 100 % of these mice against a challenge with CEA-expressing colon carcinoma cells (34). Both cellular and humoral responses have been reported after DNA vaccination in mice. In other studies a MUC-1 tandem repeat array was used for DNA vaccination of mice and 30% of these mice were protected from a tumor challenge

with MUC-1 transfected murine tumor cells (35). DNA vaccination may also be used to elicit immune responses against predefined peptide epitopes, several groups now exploit the string-bead approach to link multiple different CTL or helper epitopes together on the DNA level (36). In some cases the string-bead of peptide coding DNA is built into a vaccinia virus as a delivery vehicle. Recently, it was shown that such a vaccinia virus recombinant poly-epitope vaccine was able to protect mice against several virus infections and a tumor challenge (37). The authors show that all 10 minimal peptide epitopes encoded by the string-bead are expressed and recognized by the appropriate T cell clones (38). RNA was also shown to confer anti-tumor immunity. Vaccination with RNA to ovalbumin induced CTL in mice (39). In conclusion, multiple studies have shown the efficacy of DNA vaccines in anti-viral and anti-tumor immunity.

Thus, according to yet another aspect of the present invention there is provided a DNA vaccine composition which includes at least one polynucleotide encoding a tumor associated antigen peptide derived from a sequence encoding a protein selected from the group consisting of Uroplakin (UP), Prostate specific antigen (PSA), Prostate specific membrane antigen (PSMA), Prostate acid phosphatase (PAP), Lactadherin (BA-46), Mucin (MUC1) and Teratocarcinoma-derived growth factor (CRIPTO-1). As further detailed herein, the at least one polynucleotide can be a part of a longer polynucleotide designed to encode a fused protein product from which the tumor associated antigen peptide is cleavable by a protease.

The polynucleotide is preferably DNA in a form of, or contained in, for example, naked DNA, plasmid, retroviral vector, adenoviral vector, vaccinia viral vector, herpes viral vector, lenti virus vector, EBV vector, CMV vector, polio virus vector, sindbis viral vector, semliki forest virus vector, parvo virus vector, adeno-associated virus vector, virus like particle (VLP) vector. Alternatively, the polynucleotide can be in the form of RNA. The polynucleotide can also be delivered in a non-viral delivery system, such as, for example, but not limited to, in liposomes, in complex with cationic reagents, or with a polycation, such as poly-lysine. The polynucleotide can also be delivered by mechanical means, such as, but not limited to, a gene-gun, by electrical means, or in bacterial vectors like BCG.

There is increasing evidence that peptide vaccination may be much more effective when the peptides are introduced together with an antigen presenting cell (APC) (40). In previous studies of a murine lung carcinoma we have shown that vaccination with a defined TAA peptide (MUT-1) loaded on APC result in long term survival of mice bearing lung metastases (41, 42). The most common cells

used to load antigens are bone marrow and peripheral blood derived dendritic cells (DC), as these cells express costimulatory molecules that help activation of CTL. Preliminary clinical trials have been performed. In one trial HLA-A1 melanoma patients have been treated with autologous DC loaded with a MAGE-1 peptide. CTL activity was increased in tumor infiltrated lymphocytes (43). In another study, five patients with advanced pancreatic carcinoma were treated with a K-ras derived peptide loaded on DC. As a mutation of K-ras at codon 12 is frequently found in pancreatic carcinoma, three differently mutated peptides, 12-Asp, 12-Arg and 12-Val (non mutated sequence is 12-Gly) were used for vaccination, matched to the mutation in the patient's tumor. Two of the patients showed a specific CTL response and prolonged survival (44). A phase I clinical trial in 51 prostate cancer patients compared a soluble peptide to a DC based peptide in HLA-A2 patients. The peptide was derived from PSMA (SEQ ID NO:29). Only 7 patients that received DC based vaccines with this peptide responded by decreased levels of serum PSA (45). In animal studies a number of groups showed that macrophages loaded with peptides constitute efficient vaccines, yet the number of cells used for vaccination is 10 fold higher than equivalent DC vaccines. Recently in a murine lung carcinoma model the efficacy was tested of syngeneic fibroblasts treated with a proteasome inhibitor to decrease levels of endogenous peptides and loaded with synthetic MUT peptides as vaccines. Effective protection was found against metastatic spread of lung carcinoma.

Thus, according to still another aspect of the present invention there is provided a cellular vaccine composition which includes an antigen presenting cell presenting at least one tumor associated antigen peptide derived from a protein selected from the group consisting of Uroplakin (UP), Prostate specific antigen (PSA), Prostate specific membrane antigen (PSMA), Prostate acid phosphatase (PAP), Lactadherin (BA-46), Mucin (MUC1) and Teratocarcinoma-derived growth factor (CRIPTO-1). As further detailed herein, the antigen presenting cell can, for example, be a dendritic cell, a macrophage, a B cell and a fibroblast. Presenting the at least one tumor associated antigen peptide can be effected by a method selected from the group consisting of (a) transducing the antigen presenting cell with at least one polynucleotide (e.g., DNA) encoding the at least one tumor associated antigen peptide; (b) loading the antigen presenting cell with at least one polynucleotide (e.g., RNA) encoding the at least one tumor associated antigen peptide; (c) loading the antigen presenting cell with the at least one tumor associated antigen peptide (e.g., synthetic); and (d) loading the antigen presenting cell with at least one longer polypeptide (e.g., purified) including the at least one tumor associated antigen peptide. Loading can be external or internal. The

polynucleotide, peptide or longer polypeptide can be fused to internalizing sequences, antennapedia sequences or toxoid sequences or to helper sequences, such as, but not limited to, heat shock protein sequences.

While it is clear that CD8+ class-I restricted CTL recognize and destroy tumor cells in vitro and in vivo, animal models often show a requirement of CD4+ MHC-class-II restricted T cell help for optimal responses (83). Helper T cell epitopes can contribute to induction of cellular immune responses by class I peptide vaccines, as seen by the synergistic tumor protection upon simultaneous vaccination with T helper and CTL epitopes (84). The 'help' to CTL is most often provided via the production of specific cytokines. Helper epitopes can be specific and derived from a tumor antigen (85). They can also broadly crossreact with a number of MHC class II molecules, and may be either pathogen-derived or comprised of sequences not found in nature (86-88). More specifically, a sequence containing a T helper epitope can be linked to a CTL epitope to create one immunogenic entity. Alternatively, a mixture of 2 or more separate entities, corresponding to CTL and T helper epitopes can be administered to elicit the desired CTL response. T helper epitopes can also be conjugated to other molecules or compounds which increase their biological activity.

As used herein in the specification and in the claims section below the phrase "tumor associated antigen" also refers to tumor specific antigen.

As used herein in the specification and in the claims section below the term "peptide" refers to native peptides (either degradation products or synthetically synthesized peptides) and further to peptidomimetics, such as peptoids and semipeptoids which are peptide analogs, which may have, for example, modifications rendering the peptides more stable while in a body, or more immunogenic. Such modifications include, but are not limited to, cyclization, N terminus modification, C terminus modification, peptide bond modification, including, but not limited to, CH<sub>2</sub>-NH, CH<sub>2</sub>-S, CH<sub>2</sub>-S=O, O=C-NH, CH<sub>2</sub>-O, CH<sub>2</sub>-CH<sub>2</sub>, S=C-NH, CH=CH or CF=CH, backbone modification and residue modification. Methods for preparing peptidomimetic compounds are well known in the art and are specified in Quantitative Drug Design, C.A. Ramsden Gd., Chapter 17.2, F. Choplin Pergamon Press (1992), which is incorporated by reference as if fully set forth herein. Further detail in this respect is provided herein.


As used herein in the specification and in the claims section below the term "amino acid" is understood to include the 20 naturally occurring amino acids; those amino acids often modified post-translationally *in vivo*, including for example hydroxyproline, phosphoserine and phosphothreonine; and other unusual

amino acids including, but not limited to, 2-aminoadipic acid, hydroxylysine, isodesmosine, nor-valine, nor-leucine and ornithine. Furthermore, the term "amino acid" includes both D- and L-amino acids. Further elaboration of the possible amino acids usable according to the present invention and examples of non-natural amino acids useful in MHC-1 recognizable peptide antigens are given herein.

Peptides can be of 8, 9 or 10 amino acids and peptides of 9 or 10 amino acid residues may be desirable, preferably 9. Thus, assume the following positions (P1-P9) in a 9-mer peptide:

P1-P2-P3-P4-P5-P6-P7-P8-P9

The P2 and P9 positions include the anchor residues which are the main residues participating in binding to MHC class 1 molecules, more specifically HLA-A2. Amino acid residues engaging positions P2 and P9 are hydrophobic or hydrophilic natural amino acids or non-natural amino acids. Examples of natural amino acids being Ala, Cys, Gln, Glu, Ile, Leu, Met, Ser, Thr and Val. These residues may preferably be neutral, hydrophobic, aliphatic and more preferably Val, Leu and Ile. Examples of non-natural amino acids being norleucine (Nle), norvaline (Nva), aminobutyric acid preferably  $\alpha$ -aminobutyric acid. These residues may preferably be non charged and more preferably aliphatic. P9 can also be an aliphatic amino acid of the general formula  $\text{-HN(CH}_2\text{)}_n\text{COOH}$ , wherein  $n = 2-5$ , as well as by branched derivatives thereof, such as, but not limited to,



wherein R is, for example, methyl, ethyl or propyl, located at any one or more of the n carbons.

Positions P1 and P3 are also known to include amino acid residues which participate or assist in binding to MHC molecules, however, these positions can include any amino acids, natural or non-natural.

The amino terminal residue (position P1) can also be positively charged aliphatic carboxylic acids, such as, but not limited to,  $\text{H}_2\text{N(CH}_2\text{)}_n\text{COOH}$ , wherein  $n = 2-5$  and  $\text{H}_2\text{N-C(=NH)-NH(CH}_2\text{)}_n\text{COOH}$ , wherein  $n = 2-4$ , hydroxy Lysine,  $\text{N}^\epsilon$ -methyl Lysine,  $\text{N}^\epsilon$ -ethyl Lysine,  $\text{N}^\epsilon$ -propyl Lysine or ornithine (Orn). Additionally, the amino terminal residue can be aromatic residues, such as, but not limited to, phenyl glycine, p-aminophenyl alanine, p-guanidinophenyl alanine or pyridinoalanine (Pal). These latter residues may form hydrogen bonding with the

OH<sup>-</sup> moieties of the Tyrosine residues at the MHC-1 N-terminal binding pocket, as well as to create, at the same time aromatic-aromatic interactions

The other positions P4-P8 are engaged by amino acid residues which typically do not participate in binding to MHC molecules, rather these amino acids are presented to the immune cells. Further details relating to the binding of peptides to MHC molecules can be found in reference 82, see Table V thereof, in particular.

Amino acid residues engaging position P4-P8 can include any amino acids natural or non natural. These residues may optionally be phosphorylated and/or glycosylated. Indeed residues which have been phosphorylated or glycosylated have been shown in some cases to enhance the binding to the T cell receptor.

Cyclization can engage position P4-P8, preferably positions P6 and P7. Cyclization can be obtained through amide bond formation, e.g., by incorporating Glu, Asp, Lys, Orn, di-amino butyric (Dab) acid, di-aminopropionic (Dap) acid at various positions in the chain (-CO-NH or -NH-CO bonds). Backbone to backbone cyclization can also be obtained through incorporation of modified amino acids of the formulas  $H-N((CH_2)_n-COOH)-C(R)H-COOH$  or  $H-N((CH_2)_n-NH_2)-C(R)H-COOH$ , wherein  $n = 1-4$ , and further wherein R is any natural or non-natural side chain of an amino acid.

Cyclization via formation of S-S bonds through incorporation of two Cys residues is also possible. Additional side-chain to side chain cyclization can be obtained via formation of an interaction bond of the formula  $-(-CH_2-)_n-S-CH_2-C(=O)-$ , wherein  $n = 1$  or  $2$ , which is possible, for example, through incorporation of Cys or homoCys and reaction of its free SH group with, e.g., bromoacetylated Lys, Orn, Dab or Dap.

In longer peptides such as in a 10 mer peptide in which the second anchor amino acid is at position P10 the amino acid engaging P9 may include most L amino acids. In some cases shorter peptides such as an 8 mer peptide are also applicable, in which the carboxy terminal acid P8 may serve as the second anchor residue. All the options described for the anchor amino acid residues engaging positions P2 and P9 in a 9 mer peptide may apply likewise to the anchor amino acid residues engaging positions P2 and P10 in a 10 mer peptide and P2 and P8 in an 8 mer peptide.

The amino acids may be modified as is necessary to provide certain characteristics such as greater immunogenicity, more stability or improved pharmacological properties. The peptides can be for instance subject to changes such as the replacement of one or more amino acid residues whether dissimilar or similar.



Modification of the peptides may also be by decreasing, e.g. in a 10 mer peptide, or extending, e.g. in an 8 mer peptide, the amino acid sequence, for example, by deletion or addition of amino acids. It will be appreciated that preferably anchor amino acids should not be deleted.

5 Peptide bonds (-CO-NH-) within the peptide may be replaced by N-alkylated bonds such as N-methylated (-N(CH<sub>3</sub>)-CO-), ester bonds (-C(R)H-C-O-O-CH(R)-N-), ketomethylen bonds (-CO-CH<sub>2</sub>-), -aza bonds (-NH-N(R)-CO-), wherein R is hydrogen or any alkyl, e.g., methyl carba bonds (-CH<sub>2</sub>-NH-), hydroxyethylene bonds (-CH(OH)-CH<sub>2</sub>-), thioamide bonds  
10 (-CS-NH-), olefinic double bonds (-CH=CH-), retro amide bonds (-NH-CO-), and peptide derivatives (-N(R)-CH<sub>2</sub>-CO-), naturally presented on the carbon atom.

These modifications can occur at any of the bonds along the peptide chain and even at several (2-3) at the same time. Preferably, but not in all cases necessary, these modifications should exclude anchor amino acids.

15 For amino acid residues engaging positions other than the second residue from the amino terminal and the end residue at the carboxy terminal natural aromatic amino acids, Trp, Tyr and Phe, may be replaced by synthetic non-natural acid such as TIC, naphthylalanine (Nal), ring-methylated derivatives of Phe, halogenated derivatives of Phe or o-methyl-Tyr.

20 As used herein in the specification and in the claims section below, the term "transduced" refers to the result of a process of inserting nucleic acids into cells. The insertion may, for example, be effected by transformation, viral infection, injection, transfection, gene bombardment, electroporation or any other means effective in introducing nucleic acids into cells. Following transduction the  
25 nucleic acid is either integrated in all or part, to the cell's genome (DNA), or remains external to the cell's genome, thereby providing stably transduced or transiently transduced cells.

As used herein in the specification and in the claims section below the phrase "derived from a protein" refers to peptides derived from the specified  
30 protein or proteins and further to homologous peptides derived from equivalent regions of proteins homologous to the specified proteins of the same or other species, provided that these peptides are effective as anti-tumor vaccines. The term further relates to permissible amino acid alterations and peptidomimetics designed based on the amino acid sequence of the specified proteins or their  
35 homologous proteins.

As used herein in the specification and in the claims section below the phrase "anti-tumor vaccines" refers to vaccines effective in preventing the development of, or curing, cancer, including primary tumor and/or metastases.

As used herein in the specification and in the claims section below the phrase "prevention or cure" also refers to inhibiting, slowing or reversing the progression of a disease, substantially ameliorating clinical symptoms of a disease or substantially preventing the appearance of clinical symptoms of a disease.

5 In the specification and in the claims section below the phrase as used herein 'loading' refers to exposing, adding or introducing a substance into or onto a cell or vesicle or part thereof.

According to a preferred embodiment of the present invention the Uroplakin (UP), Prostate specific antigen (PSA), Prostate specific membrane  
10 antigen (PSMA), Prostate acid phosphatase (PAP), Lactadherin (BA-46), Mucin (MUC1) and Teratocarcinoma-derived growth factor (CRIPTO-1) proteins are each independently of a mammal, e.g., a humanoid such as a human being or a rodent such as murine.

According to yet another preferred embodiment of the present invention the  
15 composition further comprising a carrier. Usually the tumor associated antigen peptide(s) are presented in context of the carrier.

The carrier can be a proteinaceous carrier to which the peptides are linked. Methods of linking short peptides to carriers are well known in the art of vaccination. The carrier can alternatively be a particulate adjuvant, an oil or  
20 emulsifier based adjuvant, a gel based type adjuvant, or an adjuvant based on specific targeting of antigen, such as, but not limited to, antibody-liposom conjugates. The carrier can also be a protein or a recombinant protein produced, for example in bacteria, yeast or in mammalian cells, including cytokines, such as  
25 IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-14, IL-15, IL-16, IL-17, IL-18, interferon alpha, interferon beta, interferon gamma and others. The carrier can also be an antigen presenting cell, such as, but not limited to, a dendritic cell, a macrophage, a B cell or a fibroblast. The cell selected is either an autologous or non-autologous HLA matching cell. Optionally the cell can be a cultured cell, a cell treated by various reagents (e.g., by early  
30 and/or late acting cytokines), transduced by genes, and/or irradiated or radiated.

The vaccine composition according to the present invention is effective in prevention or cure of cancer and/or cancer metastases. In other words, the composition is effective for primary tumors, secondary tumors and metastases thereof in the same organ or in another organ, provided that the tumor expresses  
35 the above listed tumor associated proteins. According to a preferred embodiment of the present invention the cancer being treated or prevented via the administration of the vaccine composition is a carcinoma. i.e., a malignant tumor composed of epithelial tissue. The cancer treated or prevented according to the

present invention can be, for example, breast, bladder, prostate, pancreas, ovary, thyroid, melanoma, colon, stomach and/or head and neck cancer.

According to an embodiment of the present invention, the Uroplakin protein can be Uroplakin II, Uroplakin Ia, Uroplakin III and Uroplakin Ib.

5 According to another embodiment of the present invention, the tumor associated antigen peptides derived from the Mucin protein are from a non-tandem repeat array of Mucin.

10 According to yet another embodiment of the present invention, one or more tumor associated antigen peptides derived from the Mucin protein are from a region selected from the group consisting of a signal peptide, a cytoplasmic domain and an extracellular domain of Mucin.

According to a presently preferred embodiment of the present invention the one or more tumor associated antigen peptides are selected from SEQ ID NOs: 1 to 64 and 66 to 77 and effective homologues and analogs thereof.

15 For therapeutic or prophylactic anti-tumor treatment, the vaccine composition according to the present invention may include thickeners, carriers, buffers, diluents, surface active agents, preservatives, and the like, all as well known in the art. The composition may also include one or more active ingredients, such as, but not limited to, anti-inflammatory agents, anti-microbial agents, anesthetics and the like.

20 The vaccine composition may be administered in either one or more ways. Administration may be effected topically (including ophthalmically, vaginally, rectally, intranasally), orally, by inhalation, or parenterally, for example by intravenous drip or intraperitoneal, intravesical, subcutaneous, or intramuscular injection.

25 Compositions for topical administration can include, but are not limited to, lotions, ointments, gels, creams, suppositories, drops, liquids, sprays and powders. Conventional pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like may be necessary or desirable.

30 Formulations for parenteral administration can include, but are not limited to, sterile aqueous solutions which may also contain buffers, diluents, adjuvant and other suitable additives. The adjuvant is preferably of a type allowed for use in treating human beings, such as BCG adjuvant.

35 Dosing is dependent on responsiveness, but will normally be one or more doses per week or month, with course of treatment lasting from several weeks to several months. Persons ordinarily skilled in the art can easily determine optimum dosages, dosing methodologies and repetition rates.

The present invention provides novel tumor associated antigen peptides effective in eliciting CTL response which can therefore be effective therapeutic agents to combat cancer.

According to another embodiment of the present invention there is provided a peptide derived from a protein selected from the group consisting of Uroplakin (UP), Prostate specific antigen (PSA), Prostate specific membrane antigen (PSMA), Prostate acid phosphatase (PAP), Lactadherin (BA-46) and Mucin (MUC1) and Teratocarcinoma-derived growth factor (CRIPTO-1), the peptide comprising 8-10 amino acid residues as selected so as to promote effective binding to a MHC class 1 type molecule such that a CTL response is elicitable

Each of the various aspects of the present invention as delineated hereinabove and as claimed in the claims section below finds experimental support in the Examples section that follows.

## EXAMPLES

Reference is now made to the following examples, which together with the above descriptions, illustrate the invention in a non limiting fashion.

## MATERIALS AND EXPERIMENTAL METHODS

**Mice:** The derivation of HLA-A2.1/D<sup>b</sup>- $\beta$ 2m monochain, transgenics, H-2D<sup>b</sup> x  $\beta$ 2m double knockout mice (HhD mice) was described before (30).

**Tumor and cell lines:** RMA-S is a TAP-2 deficient lymphoma clone of C57BL/6 origin. The RMA-S HhD and RMA-S-HhD-B7.1 are transfectants carrying the HhD construct without or with the gene for murine B7-1 costimulatory molecule, respectively. T2 is a TAP-2 deficient human lymphoblastoid line expressing HLA-A2. RMA-S, RMA-S derivatives and T2 cells were maintained in RPMI 1640-10 % FCS and antibiotics.

The human tumor cell lines, J82 (TCC of the bladder) sup (TCC of the bladder, DU145 (prostate carcinoma) and MDA-MB-157 (breast carcinoma) and the HhD transfectants of DU145 and MDA-MB-157 were maintained in DMEM-10 % FCS 2 mM glutamine - 1 mM sodium pyruvate - 1 % non-essential amino acids and antibiotics.

HhD and B7-1 transfectants were grown in presence of 500-1000  $\mu$ g/ml of Genitcin, G418. All tissue culture media and supplements were purchased from Gibco.

**Peptide synthesis:** Peptides were synthesized on an ABIMED AMS 422 multiple peptide synthesizer (Langenfeld, Germany), employing the a-N-9-fluorenylmethoxycarbonyl (Fmoc) strategy following the company's commercially available protocols. Peptide chain assembly was conducted on a 2-chlorotrityl chloride resin (Novabiochem, Laufelfingen, Switzerland). Crude peptides were purified to homogeneity by reversed-phase HPLC on a semi-preparative silica C-8 column (250 x 10 mm; Lichnonorb RP-8; Merck). Elution was accomplished by a linear gradient established between 0.1 % trifluoroacetic acid in water and 0.1 % trifluoroacetic acid in 70 % acetonitrile in water (v/v). Purity of peptides ( 96 %) was ascertained by analytical HPLC (RP-18) using various acetonitrile water (containing 0.1 % trifluoroacetic acid) gradients. The products' compositions were determined by amino acid analysis (Dionex automatic amino acid analyzer, Sunnyvale, CA) after extraction acid hydrolysis. Molecular weight was ascertained by mass spectrometry (VG Tofspec; Laser Desorption Mass. Spectrometry; Fison Instruments, Manchester, U.K.). Tables 1-8 present these peptides.

**Preparation of peptide fractions from fresh human tumors and normal tissues:** Surgical specimens of TCC, normal bladder mucosa, prostate carcinoma, hyperplastic benign prostate, breast carcinoma or normal breast tissue were transferred on ice from surgery to a qualified pathologist. Tumor tissue were cleaned from most adjacent tissue and snap frozen. Non-necrotic fragments, from 5-10 patients were homogenized in PBS; 0.5 % Nonidet P-40; 10 µg/ml soybean trypsin inhibitor; 5 µg/ml lupeptide; 8 µg/ml aprotinin and 0.5 mM PMSF and homogenized using a glass-teflon homogenizer. Following further stirring for 30 minutes at 4°C, the homogenates were titrated with 10 % TFA to a final concentration of 0.1 % TFA and stirred for 30 minutes at 4°C. After ultra-centrifugation for 30 minutes at 42K rpm the supernatants were applied to Sephadex G25 columns and fractions were monitored at 230 nm. Peptide fractions under 10 Kd were pooled, lyophilized and further fractionated by centriprep 3 by centrifugation (Amicon, Beverly, MA). The peptide pool after lyophilization, was dissolved in opti-MEM (Gibco) for further use.

**Scoring of HLA-A2.1 binding peptides:** Protein sequences were screened for MHC binding by a HLA Peptide Binding Predictions software approachable through a worldwide web interface at (see also reference 82). This software, based on accumulated data, scores every possible peptide in the protein for possible binding to MHC according to the contribution of every amino acid in the peptide. Theoretical binding scores represent calculated half-life of the HLA-A2.1-peptide complex.

**Measurement of peptide binding by stabilization of cell surface MHC:**

Peptide binding to HhD was measured by stabilization of HhD on RMA-S transfectants, using an indirect FACS assay as follows:  $5 \times 10^5$  peptide loaded TAP-2 deficient RMA-S-HhD cells (see vaccination), were incubated with anti HLA monoclonal antibodies for 30 minutes at 4°C. After washing the cells with PBS - 0.5 %, BSA + 0.1 %, sodium azide, the second antibody, goat anti mouse-FITC (Jackson Lab.), was applied for 30 minutes at 4°C. Following washing, the amount of bound antibodies was detected by FACSScan (Bacson).

Mouse monoclonal antibodies B-9-12, w6/32 (anti HLA A, B, C) and B.B7.2 (anti HLA-A2.1) were used for analyses.

**Vaccination:** Mice were immunized intraperitoneally three times at 7-day intervals, with  $2 \times 10^6$  irradiated (5,000 rad) tumor cells, or with peptide-loaded RMA-S-HhD-B7.1 transfectants. Peptide loading of RMA-S-HhD-B7.1 cells was performed as follows: After washing the cells 3 times in PBS, the surface expression of HhD monochins was stabilized by a 4 hours culture at 26°C. Synthetic peptides, or peptide extracts were added to  $10 \times 10^6$  cells in 1 ml of opti-MEM (Gibco) medium to a concentration of 100  $\mu$ M or 1 mM, respectively. The cells were incubated at 26°C and for additional 3 hours at 37°C. Peptide loaded RMA-S-HhD-B7.1 cells were irradiated (5000 rad), washed, resuspended in PBS and injected into mice. In mixed synthetic vaccines, RMA-S-HhD-B7.1 cells were loaded separately with each peptide and pooled before vaccination.

**In vitro cytotoxicity assays:** Mice were immunized intraperitoneally three times at 7-day intervals, with  $2 \times 10^6$  irradiated (5,000 rad) tumor cells, or with peptide-loaded RMA-S-HhD-B7.1 transfectants. Spleens were removed on day 10 after the last immunization and splenocytes were restimulated *in vitro* with either irradiated tumor cells (for mice immunized with tumor cells) or with lymphocytes pulsed with 100  $\mu$ M synthetic peptides or 1 mM patients-derived extract in opti-MEM (Gibco) medium. Restimulated lymphocytes were maintained in RPMI 1640 medium containing 10 % FCS, 1 mM glutamine, combined antibiotics, 1 mM sodium pyruvate, 10 mM HEPES, pH 7.4,  $5 \times 10^{-5}$  M  $\beta$ -mercaptoethanol, and 1 % nonessential amino acids (RPMI-HEPES medium) for 5 days. Viable lymphocytes (effector cells), were separated by lymphocyte-M (Cedarlane, Ontario, Canada) centrifugation, resuspended in RPMI-HEPES medium, and admixed at different ratios with  $5 \times 10^3$  peptide loaded RMA-S-HhD cells, previously loaded with different peptides (see vaccination). CTL assays were performed in U-shaped microtiter wells, at 37°C, 5 % CO<sub>2</sub> for 5 hours. Cultures were terminated by centrifugation at 1,000 rpm for 10 minutes at 4°C. A

total of 100  $\mu$ l of the supernatants was mixed with scintillation fluids and counted in a beta counter. Percentage of specific lysis was calculated as follows: % lysis = (cpm in experimental well - cpm spontaneous release)/(cpm maximal release - cpm spontaneous release) x 100. Spontaneous release was determined by incubation of 100  $\mu$ l-labeled target cells with 100  $\mu$ l of medium. Maximal release was determined by solubilization of target cells in 100  $\mu$ l 0.1 M NaOH.

**Sensitization of human PBL *in vitro*:** Patients were HLA typed by the Tissue Typing Unit of The Rabin Medical Center, Israel. Only HLA-A2.1 carrying haplotypes were used herein. Heparanized blood, 40 ml/patient was separated on Ficoll gradients, mononuclear cells were collected at the interphase, washed three times in PBS and resuspended in PBS. A third of the lymphocyte volume was incubated with tumor extracted peptides, at a concentration of 1 O.D230/ml for 2 hours at 37°C. The other two thirds of the cells were suspended at 10<sup>6</sup> cells/ml in RPMI - 10 % FCS, 2 mM glutamine, 1 mM sodium pyruvate, 10 mM Hepes, 5 x 10<sup>-3</sup> M  $\beta$ -mercaptoethanol and combined antibiotics. Peptide loaded cells were added together with 5-20 U/ml of recombinant human IL-2 (Peptotec, Rehovot, Israel). Cells were incubated for 5-7 days at 37 °C, 5 % CO<sub>2</sub> and then used in CTL assays as described above.

## EXPERIMENTAL RESULTS

### EXAMPLE 1

#### Uroplakins are T cell defined TAAs in transitional cell carcinoma (TCC)

Transitional cell carcinoma (TCC) of the bladder represents a prevalent cancer. Its incidence increases gradually during life and peaks in the 7th and 8th decade to 200/100,000. About 95 % of bladder tumors are TCC. Of these, 75 % are superficial papillary tumors and have a 15 % - 20 % progression rate during the first two years of follow up. Approximately, 20 % - 25 % of TCC present as muscle invasive tumors with aggressive behavior and a 40 % survival rate at 5 years (46). After surgical resection of superficial tumors, intravesical instillations of chemotherapeutics as Tiotepa or Mitomycin, or non-specific intravesical immunotherapy with Bacillus Calmette Guerein (BCG), reduce the frequency of recurrences and have a minimal effect on progression. BCG intravesical instillation was found to be the most effective in reducing the recurrence and progression rate of high grade papillary TCC and of transitional cell carcinoma *in situ*. The mechanism of action was not yet elucidated. However, following therapy there is a mononuclear infiltrate of the lamina propria of the bladder

mucosa. This infiltrate is rich in T-cells, macrophages and Langerhans cells (47). It was found that CD8 T-cells were activated after BCG intravesical therapy and induced specific lysis of TCC cells (48). Also, BCG intravesical instillation resulted in increased presentation of MHC complexes on transitional cell membrane (49). Lately, CTL, which induced specific lysis of autologous tumor cells, were identified among tumor infiltrating lymphocytes grown from transitional cell carcinomas of the bladder (50). It seems that specific cytotoxic T-cell immunity against TCC may be augmented by BCG intravesical therapy.

An important group of TAAs are actually differentiation antigens. Such antigens were shown to induce CTL in melanoma and in colon carcinoma patients (22, 23).

Transitional epithelium is a differentiated multilayered epithelium which is restricted to the urinary tract. It is composed of 4 to 7 cell layers: a basal cell layer; an intermediate 2-3 cell layer and an apical (luminal) layer composed of umbrella like cells. These latter cells are connected by tight junctions and create an impermeable barrier to water and urinary solutes.

Lately, differentiation proteins of transitional epithelium have been characterized morphologically and biochemically. These proteins were found in urothelial plaques which are present in the luminal plasma membrane of urothelial superficial (umbrella) cells. The molecular constituents of these plaques comprise four transmembranal proteins designated Uroplakins (UP); UP Ia (27 Kd); UP Ib (28 Kd); UP II (15 Kd) and UP III (47 Kd glycoprotein; the size of the core protein is 28.9 Kd). Uroplakin III probably plays a role in the formation of the urothelial glycocalyx and may interact with the cytoskeleton. An extensive evaluation of UP tissue distribution showed that UP are restricted to normal transitional cell epithelium and to TCC (51). In normal transitional epithelium the UPs were found in the apical membrane of the superficial umbrella cells. In superficial papillary TCC, the UPs were localized in the apical membrane of the luminal cells and in the cell membranes lining intra and inter-cellular lumina. In invasive TCC, UPs were localized more randomly in the cell membrane of the cells which infiltrated the stroma. Recently, the sequence of human Uroplakin II cDNA and Uroplakin Ia genomic sequence were determined and their protein sequences were deduced (NCBI accession Nos. 2190407 and 2098577, respectively). Since TCC patients treated by BCG were shown to have increased T cell infiltrates in the bladder mucosa, we first tested whether BCG treated patients show increased CTL activity against TCC derived peptides and whether UP II peptides may constitute CTL specific epitopes. We further used the TCC



model to test the T cell repertoire overlap between patient derived CTL and CTL induced in the unique H-2D<sup>b</sup><sup>-/-</sup> x  $\beta$ 2m<sup>-/-</sup> x HhD<sup>+/+</sup> (HhD) mice.

**Cytotoxic activity of human peripheral blood lymphocytes (PBL) against target cells pulsed with Uroplakin II peptide homologues:** The human UP II amino acid sequence was screened for potential HLA-A2 binding, 9 amino acids peptides by a HLA binding motif program. We chose that particular HLA allele because it is present in 45 % of the population. Seven 9-mer peptides, which were predicted to bind with high affinity to HLA-A2 were selected and synthesized (Table 1). The binding of these peptides to HLA-A2 was tested by an MHC stabilization assay utilizing TAP negative cells, which can present exogeneously bound peptides. The RMA-S cells (30) transfected by the HhD construct were loaded with various concentrations of UP II peptides and then reacted with anti-HLA antibody. High affinity binding was shown for all 7 peptides by FACS analyses (Figure 1).

TABLE 1

*Predicted human Uroplakin II peptides that bind to HLA-A2*

	Peptide	Sequence	Amino-acids*	SEQ ID:
20	HU01	FLLVLGFII	168-176	1
	HU02	VLPSVAMFL	161-169	2
	HU03	LVLGFIIAL	170-178	3
	HU04	KVVTSSFVV	67-77	4
	HU05	LVPGTKFYI	109-117	5
25	HU06	LLPIRTLPL	4-12	6
	HU07	YLVKKGAT	119-127	7

The prediction program is described in Materials and Methods. The peptides are listed in descending order of predicted HLA-A2.1-peptide complex stability. Single letter amino acid codes and the position of the peptide in the protein are listed. \* Numbering is according to NCBI accession No. 2190407.

To restimulate CTL activity in patient's PBL we chose 6 TCC patients of the HLA-A2 haplotypes: 4 patients underwent adjuvant intravesical BCG instillations and 2 underwent no adjuvant therapy after endoscopic resection of the TCC. Another patient without TCC served as control. PBL were stimulated *in vitro* by a mixture of tumor acid extracted peptides from 9 TCC samples as described before (52). Each tumor was extracted individually and an equimolar

ratio of each sample was added to the mixture. The final concentration of peptides in the stimulation reaction was 1 O.D. 230/ml. Antigen presenting cells (APC) from the same peripheral blood were pulsed with the tumor extract. Peripheral blood lymphocytes (PBL) were stimulated with APC loaded with pooled TCC extract and with IL2 for 5 days. Uroplakin II peptides, TCC extracted peptides and normal bladder extracted peptides were loaded on labeled T2 cells (HLA-A2 positive cells which are TAP deficient, express empty MHC molecules and can be loaded only with exogenous peptides), and were screened by CTL obtained from PBL of the patients, as described. CTL from TCC patients treated with BCG induced 32 % to 71 % specific lysis of Uroplakin II peptide loaded T2 cells (Figure 2). Tumor extract loaded targets were lysed at 62% while normal bladder extract loaded targets were lysed at 29 % only, indicating preferential recognition of tumor antigens. CTL from TCC patients without BCG adjuvant therapy induced 25 % to 48 % specific lysis of T2 cells loaded Uroplakin II peptides, 29 % lysis of tumor extract loaded T2 cells and 15 % lysis of bladder extract loaded T2 cells. There was no lysis of T2 cells loaded with breast tumor extract, or with a HLA-A2 binding peptide homologue to tyrosinase, a melanoma differentiation antigen. A TCC cell line expressing HLA-A2, J82 was effectively killed while another TCC line SUP, was not lysed. Also, there was no lysis of a natural killer (NK) sensitive cell line (K562). CTL obtained from the patient without TCC induced no specific lysis of tumor extract, or Uroplakin II peptides loaded APC cells. The differences between TCC patients treated with BCG, TCC patients without adjuvant therapy and non-TCC patients were very significant ( $P < 0.001$ ). We also tested the activity of patient's PBL, resensitized by the pool of tumor extracts, against individual tumor extracts. Figure 3 shows higher activity of lymphocytes derived from BCG treated patients than lymphocytes derived from non-treated patients. Control lymphocytes show low activity. Thus shared antigens are expressed in all TCC samples.

**CTL induced in HhD mice recognize the same UP II peptides as human PBL:** The immunogenicity of Uroplakin homologue peptides was also evaluated in a mouse model, in which the murine MHC class I genes are not expressed ( $D^b^{-/-}$ ,  $\beta 2$ -microglobulin  $-/-$ , double knockout) and which is transgenic for a HLA-A2 and human  $\beta 2$ -microglobulin monochain (HhD mice). The advantage of a HLA-A2 transgenic mouse model is that CTL epitopes can be detected more easily and reproducibly, without lengthy and repeated *in vitro* stimulations. The murine MHC knockout mice used herein has the additional advantage that their CTL repertoire is only HLA-A2 restricted and the CTL response against human tumor extract or human cell lines is not masked by a

potential xenogeneic response. HhD mice were immunized with the 7 Uroplakin II homologue peptides or with the same TCC peptide extract used for stimulation of patient's PBL. Lymphocytes were obtained from the spleens of these mice and restimulated *in vitro* once with the corresponding peptides.

5 The 7 UP II peptides, that were recognized by human CTL, 8 new UP Ia homologues peptides with high HLA-A2 binding affinity (Table 2), 4 murine UP II derived HLA-A2 binding peptides (Table 3), TCC and normal bladder mucosa peptide extracts and two TCC cell lines: J82 (HLA-A2 positive) and TCCSUP were screened by the CTL obtained from these mice.

10

TABLE 2

*Predicted human Uroplakin Ia peptides that bind to HLA-A2*

Peptide	Sequence	Amino-acids*	SEQ ID:
HUO11	SLFAETIWV	33-41	8
15 HUO12	MLIAMYFYT	248-256	9
HUO13	LMWTLPVML	241-249	10
HUO14	MLIVYIFEC	100-108	11
HUO15	YIFECASCI	104-112	12
HUO16	LVLMLIVYI	97-105	13
20 HUO17	ALCRRRSMV	85-93	14
HUO18	LLSGLSLFA	28-36	15

The prediction program is described in Materials and Methods. The peptides are listed in descending order of predicted HLA-A2.1-peptide complex stability. Single letter amino acid codes and the position of the peptide in the protein are listed. \* Numbering is according to NCBI accession No. 2098577.

25

TABLE 3

*Predicted murine Uroplakin II peptides that bind to HLA-A2*

Peptide	Sequence	Amino-acids*	SEQ ID:
30 MUO1	FLLVVGLIV	168-176	16
MUO3	LVVGLIVAL	170-178	17
MUO4	KVVKSDFFV	69-77	18
MUO6	TLPVQTLPL	4-12	19

The prediction program is described in Materials and Methods. The peptides are listed in descending order of predicted HLA-A2.1-peptide complex stability. Single letter amino acid codes and the position of the peptide in the protein are listed. The murine 1, 3, 4, 6 peptides are homologous to the human Uroplakin II

35

peptides 1, 3, 4, 6 in Table 1. \* Numbering is according to NCBI accession No. 586160.

The results are expressed as lytic units (LU30), which represent  $10^6$ /the number of effectors which induce 30 % specific target lysis. More LU signify a more potent CTL response. Figure 4 shows that CTL from HhD mice immunized with Uroplakin II peptides (1-7 loaded individually and injected by a mixture of loaded cells) induced a significant lysis of RMA-S-HhD cells loaded with Uroplakin II homologue peptides (8.1 to 32.2 LU). There was no lysis of RMA-S-HhD cells loaded with nonspecific peptides as breast tumor extract or the HLA-A2 binding tyrosinase homologue peptide. Also, there was no lysis of RMA-S-HhD cells without peptides (empty target cells). There was intense lysis of J82 cells (17.3 LU) and there was much less lysis of TCCSUP cells (1.2 LU). There was also intense lysis of RMA-S-HhD cells loaded with TCC peptide extract (15 LU) and less lysis of normal bladder mucosa peptide extract loaded cells (4.5 LU) (Figure 4).

CTL from mice immunized with TCC peptide extract induced significant lysis of RMA-S-HhD targets loaded with Uroplakin II homologue peptides (3.2 to 14 LU); Uroplakin Ia homologue peptides (2 to 11.4 LU); TCC peptide extract (12.6 LU) and less lysis of these targets loaded with normal bladder mucosa peptide extract (2 LU). There was also intense lysis of J82 cells (7.6 LU) and less lysis of TCCSUP cells (0.8 LU). There was no lysis of target cells loaded with nonspecific peptides or of empty target cells. The lysis pattern of the targets loaded with the different Uroplakin II homologue peptides was similar to the CTL assays using human PBL, CTL from HhD mice immunized with Uroplakin II peptides or with TCC peptide extract (Figure 5).

Although human and HhD murine CTL recognize the same UP II peptides, it was not clear whether HhD murine CTL are directed against the actual T cell epitopes or against possible differences between humanoid and rodent UP II. We evaluated also the crossreactivity of CTL from mice immunized with human Uroplakin II peptides against 4 homologue murine Uroplakin II peptides. The murine UP II sequence was screened for HLA-A2 binding peptides and 4 peptide homologues to human peptides 1, 3, 4 and 6 were synthesized. Murine peptides differ in 1-3 amino acids from similar human peptides, yet most are conservative changes (Tables 1 and 3). Murine UP II peptides bind stably to RMA-S-HhD cells.

There was significant lysis of the targets loaded with each of these murine peptides. A good correlation between the LU30 results for the human and their

homologue murine peptides was found. Three human peptides induced more target lysis than their murine counterpart and for the fourth peptide the trend was opposite (Figures 4 and 5). Thus, the specificity of the HhD murine CTL, as the specificity of human PBL is directed to common epitopes on the peptides. To  
5 examine whether any autoimmune effects are observed in peptide vaccinated HhD mice, the internal organs of the HhD mice immunized with TCC peptide extract or with Uroplakin II homologue peptides were formalin fixed and representative sections from each organ were stained with hematoxylin and eosin. Two non-immunized HhD mice served as control. There were no pathological finding  
10 such as tissue necrosis or inflammatory infiltrate in any of the organ examined. The urinary bladder was carefully examined using a larger number of sections. The bladder wall including the mucosa was normal in the immunized and in the control mice.

The immunogenicity of additional Uroplakin-derived peptides was  
15 evaluated in the HhD mouse strain, in which the murine MHC class I genes are not expressed (Db<sup>-/-</sup>,  $\beta$ 2-microglobulin<sup>-/-</sup> double knock-out), and which is transgenic for an HLA-A2 and human  $\beta$ 2-microglobulin monochain. Seven HLA-A2 binding peptides derived from the sequence of Uroplakin Ib, 2 from Uroplakin II, and 6 from Uroplakin III (see Table I) were tested for their ability to  
20 evoke specific CTL responses in these mice. After three weekly immunizations, splenic lymphocytes were restimulated *in vitro* with cognate peptide and then incubated with radiolabeled peptide-loaded target cells. Figures 23a-e show that several of the tested peptides (peptides B1, B2, B5, B6, 3.2, 3.3, and 8) could elicit lymphocytes which specifically lysed target cells. Importantly, in all assays there  
25 was no significant lysis of unloaded target cells (non) or of cells loaded with a non-specific HLA-A2 binding peptide from the melanoma antigen tyrosinase (tyr).

Table 3a below shows the sequences of the peptides tested in single letter amino acid code, their starting position in the intact protein.

**TABLE 3a**  
***Predicted human Uroplakin Ib, II and III peptides that bind to HLA-A2***

Peptide	Start Position	Sequence
Uroplakin Ib/B1	239	AILCWTFWV
Uroplakin Ib/B2	92	FILMFIVYA
Uroplakin Ib/B3	29	LTAECIFFV
Uroplakin Ib/B4	154	MLQDNCCGV
Uroplakin Ib/B5	240	ILCWTFWVL
Uroplakin Ib/B6	86	KILLAYFIL
Uroplakin Ib/B7	64	FVGICLFCL
Uroplakin II/8	161	VLLSVAMFL
Uroplakin II/9	162	LLSVAMFLL
Uroplakin III/3.1	214	ILGSLPFFL
Uroplakin III/3.2	128	ILNAYLVRV
Uroplakin III/3.3	221	FLLVGFAGA
Uroplakin III/3.4	20	NLQPQLASV
Uroplakin III/3.5	47	CMFDSKEAL
Uroplakin III/3.6	62	YLYVLVDSA
Tyrosinase	368	YMDGTMSQV

Table 3a shows the sequences of the peptides tested in single letter amino acid code and their starting position in the intact protein (according to NCBI accession nos. 3298345 (peptides 3.1-3.6), 3483011 (peptides 8 and 9), and 3721858 (peptides B1-B7)).

## EXAMPLE 2

**PSA, PSMA and PAP homologue peptides are immunogenic CTL epitopes in HLA-A2 transgenic H-2D<sup>b</sup>,  $\beta$ 2m double knockout mice**

Carcinoma of the prostate (CAP) is the most prevalent cancer and the second cause of death in man (53). The use of prostate specific antigen (PSA) in early detection of CAP has caused a stage shift of the disease (54). By now, 70% of CAP cases are detected when they are still organ confined. These patients may be cured by radical prostatectomy or by radiation therapy. However, the long term cancer specific survival of organ confined disease after radical prostatectomy, even in the best series, is only 70 % (55). Patients with large tumors, high grade tumors and those with seminal vesicle invasion, or positive lymph nodes are at increased risk of metastases and death due to CAP (56). However, CAP is a slow growing tumor and there is no effective adjuvant chemotherapy (57). Androgen ablation by surgical or medical means does not

increase survival, but only increases time to recurrence and palliate symptomatic bone metastases (58). There is a need for an effective adjuvant or neo-adjuvant therapy for CAP and for systemic therapy for metastatic CAP. Thus, specific immunity is being investigated as a potential adjuvant therapy. PSA and prostate specific membrane antigen (PSMA) are proteins which are expressed almost exclusively in prostate tissue, benign or malignant and are differentiation antigens. Prostate acid phosphatase (PAP) is expressed also in other tissues but its concentration is much higher in prostate tissue. In a few studies, 3 PSA homologue peptides (59-60) and 2 PSMA homologue peptides (61-62) were found to be immunogenic and to induce specific CTL by repeated *in vitro* stimulation of peripheral blood lymphocytes from CAP patients. In one of these studies (62), patients with metastatic prostate cancer were immunized with the 2 immunogenic PSMA peptides, loaded on dendritic cells. There was some partial responses expressed by a reduction in PSA levels and some objective reduction in bone metastases burden. However, the procedures used to induce CTL lines *in vitro* are cumbersome, lengthy and not reliable. Some CTL peptide epitopes which bind with high affinity to MHC class I may induce tolerance and will be not detected. Moreover, repeated *in vitro* stimulations, necessary for inducing CTL lines may select CTL which have a growth advantage in culture and not the most potent and specific CTL *in vivo*. Herein the HLA-A2 transgenic, H-2D<sup>b</sup><sup>-/-</sup>  $\beta$ 2m<sup>-/-</sup> double knockout mice (HhD mice) was used for identification of novel prostate cancer specific CTL epitopes. Peptides homologue to PSA, PSMA and PAP, which are expressed mainly in benign or malignant prostate tissue, were evaluated for their capacity to induce specific CTL in HhD mice. The amino acid sequences of these proteins (NCBI accession Nos. 296671, 2897946 and 439658) were screened by a HLA binding motif program. The peptides which were predicted to bind to HLA-A2 were synthesized (Tables 4, 5, 6) and their binding affinity was tested on RMA-S-HhD cells (TAP deficient cells, capable of presenting only exogenous peptides and which were transfected with the HhD construct) by FACS (Figures 6 and 7). The peptides with high binding affinity were used for immunization of HhD mice. Five PSA homologue peptides, in addition to the peptides detected by other authors, 6 PSMA homologue peptides, including the 2 peptides detected by other authors (PSMA 5 and 6) and 4 PAP homologue peptides were found.

TABLE 4

*Predicted human prostate specific antigen (PSA) peptides that bind to HLA-A2*

Peptide	Sequence	Amino-acids*	SEQ ID:
PSA1	DLHVISNDV	175-183	20

PSA2	VLVHPQWVL	53-61	21
PSA3	FLRPGDDSS	110-118	22
PSA4	ALGTTTCYAS	147-155	23
PSA5	KLQCVDLHV	170-178	24

- 5 The prediction program is described in Materials and Methods. The peptides are listed in descending order of predicted HLA-A2.1-peptide complex stability. Single letter amino acid codes and the position of the peptide in the protein are listed. \* Numbering is according to NCBI accession No. 296671.

10

TABLE 5

*Predicted human prostate specific membrane antigen (PSMA) peptides that bind to HLA-A2*

Peptide	Sequence	Amino-acids*	SEQ ID:
PSMA1	ELAHYDVLL	109-117	25
15 PSMA2	NLNGAGDPL	260-268	26
PSMA3	TLRVDCTPL	461-469	27
PSMA4	MMNDQLMFL	663-671	28
PSMA5	ALFDIESKV	711-719	29
PSMA6	LLHETDSAV	4-12	30

- 20 The prediction program is described in Materials and Methods. The peptides are listed in descending order of predicted HLA-A2.1-peptide complex stability. Single letter amino acid codes and the position of the peptide in the protein are listed. \* Numbering is according to NCBI accession No. 2897946.

25

TABLE 6

*Predicted human prostate acid phosphatase (PAP) peptides that bind to HLA-A2*

Peptide	Sequence	Amino-acids*	SEQ ID:
PAP1	VLAKELKFV	30-38	31
30 PAP2	ILLWQPIPV	135-143	32
PAP3	DLFGIWSKV	201-209	33
PAP4	PLERFAELV	352-360	34

- The prediction program is described in Materials and Methods. The peptides are listed in descending order of predicted HLA-A2.1-peptide complex stability.
- 35 Single letter amino acid codes and the position of the peptide in the protein are listed. \* Numbering is according to NCBI accession No. 439658.



HhD mice were immunized and boosted twice with the PSA, PSMA, or PAP peptides. Peptides were loaded individually on RMA-S-HhD cells and equal numbers of pulsed cells were mixed in the vaccine. Mice were also immunized with CAP peptide extract, or with DU145-HhD, a CAP cell line transfected with the HhD construct. Lymphocytes were obtained from the spleen of these mice and restimulated once, *in vitro* with the corresponding peptide tumor extract or cells. PSA, PSMA, PAP, CAP and normal prostate peptide extract and DU145-HhD cells were screened by the CTL obtained from these mice. The results are expressed as lytic units (LU) which represent  $10^6$ /Number of CTL effectors which induce 30 % specific target lysis. More lytic units signify a more potent CTL response.

CTL from HhD mice immunized with PAP peptides induced intense lysis of RMA-S-HhD cells (targets) loaded with PAP homologue peptides (20 to 65 LU, Figure 8). Three out of 5 PSA homologue peptides induced significant and specific lysis of targets loaded with the corresponding peptide (5.2 to 11.4 LU, Figure 9). Four out of 6 PSMA homologue peptides induced significant lysis of the targets loaded with the corresponding peptide (5 to 17 LU, Figure 10). Two out of these 4 peptides were found to be immunogenic by other authors, by *in vitro* stimulation of human peripheral blood lymphocytes. CTL obtained from each group of immunized HhD mice induced significant lysis of targets loaded with CAP peptide extract (4.5 to 22 LU). There was significantly less lysis of normal prostate peptide extract loaded targets than of the CAP peptide extract loaded targets for each immunizing peptide (Figures 8, 9 and 10). Also, there was intense lysis of DU-145-HhD cells by CTL obtained from HhD mice immunized with these cells (Figure 11), or immunized by PAP homologue peptides, or CAP peptide extract (28, 19 and 11 LU, respectively, Figure 12). There was no significant lysis of DU145-HhD cells by CTL derived from HhD mice immunized with PSA, or PSMA homologue peptides. These findings result probably from low expression of PSA and PSMA by DU145 cells.

There was good cross reactivity between CTL derived from mice immunized with CAP peptide extract and PSA, PSMA and PAP homologue peptides, which were found to be immunogenic in HhD mice. The CTL derived from mice immunized with DU145-HhD cells cross reacted with the immunogenic PAP homologue peptides (11 to 26 LU) CAP peptide extract (12 LU), and less with normal prostate peptide extract (3 LU). Overall, there was no lysis of targets loaded with nonspecific peptides such as breast cancer peptide extract, or HLA-A2 binding tyrosinase homologue peptide. Also, there was no lysis of RMA-S-HhD cells without peptides and there was no lysis of non transfected DU 145 cells

5

**EXAMPLE 3****Breast specific CTL induction by BA-46 (Lactadherin) peptides and by MUC1 peptides in HhD mice**

5 Breast cancer is the second leading cause of cancer death among women in the Western World and the leading cause of death among women at the age of 30 to 70. Breast cancer afflicts 200,000 women per annum in the USA today. The highest mortality is restricted to patients whose regional lymph nodes are involved. Early detection, followed by surgery provides good prognosis. In  
10 patients with occult lymph node metastasis, adjuvant chemotherapy or hormonal therapy for breast cancer have been proven to be effective, yet a large fraction of patients will succumb to metastasis. (63).

As pointed out above, TAA vaccines may constitute an additional treatment modality for residual disease. A number of tumor associated antigens have been  
15 described for breast carcinomas. The MUC-1 Mucin, a high molecular weight glycoprotein is highly expressed on breast carcinomas. Specific, MHC-unrestricted recognition of MUC-1 by human cytotoxic T cells were demonstrated (64). More recently peptides from MUC-1 were shown also to induce MHC class I and MHC class II restricted responses (65, 66). HER2/neu  
20 derived peptides were shown to be recognized by CTL from breast carcinoma patients (67).

BA-46 is a 46 kDa transmembrane-associated glycoprotein of the human milk fat globule membrane (HMFG), that is overexpressed in human breast carcinomas (68). The protein contains cell adhesion sequences (RGD), supports  
25 RGD based adhesion and interacts with integrin (69). It also contains an EGF-like domain, and a phospholipid binding sequence C1/C2-like domain of coagulation factor V and VIII. BA-46 is present in the circulation of breast cancer patients but not in healthy individuals (70). Moreover, anti BA-46 radio-conjugated monoclonal antibodies, have successfully targeted human breast tumors  
30 transplanted into mice (71). Herein we tested MUC1 and BA-46 derived peptides as potential TAA peptides in the HhD mouse system. BA-46 homologous peptide vaccines induce HLA-A2 restricted CTL which preferentially recognize breast tumor derived peptides.

BA-46 is overexpressed in many breast carcinomas, yet, no cellular  
35 immunity to BA-46 protein has been reported so far. We evaluated peptides complementary to the amino acid sequences of human BA-46, The sequence was screened for HLA-A2 binding motifs by a HLA binding motif program. Seven

9-mer peptides which were predicted to bind with high affinity to HLA-A2 were selected and synthesized (Table 7).

TABLE 7

5 *Predicted human breast associated BA-46 peptides that bind to HLA-A2*

Peptide	Sequence	Amino-acids*	SEQ ID:
BA-46-1	KQGNFNAWV	271-279	35
BA-46-2	NLLRRMWVT	131-139	36
BA-46-5	NLFETPILA	356-364	37
10 BA-46-6	NLFETPVEA	194-202	38
BA-46-7	GLQHWVPEL	97-105	39
BA-46-8	VQFVASVKV	313-321	40
BA-46-9	RLLAALCGA	5-13	41

15 The prediction program is described in Materials and Methods. The peptides are listed in descending order of predicted HLA-A2.1-peptide complex stability. Single letter amino acid codes and the position of the first amino acid of the peptide in the protein sequence as well as the calculated binding score are listed. \* Numbering is according to NCBI accession No. 1589428.

20 All peptides bound well to the HhD molecules expressed on RMA-S transfectant (Figure 13). HhD mice were vaccinated three times at weekly interval using  $2 \times 10^6$  RMA-S-HhD/B7.1 with either tumor extract prepared from 5 samples of breast carcinoma by acid extraction and separation of molecules < 3 Kd, or with a pool of BA-46 peptides. Ten days after the last immunization, HhD  
 25 spleen-derived antigen presenting cells (APC) were pulsed for 3 hour at 37°C with either breast tumor extract or a BA-46 peptide pool, followed by incubation with the rest of the splenocytes for 4 more days. BA-46 peptides, breast extracted peptides and normal breast extracted peptides were loaded on labeled RMA-S-HhD cells and were screened by cytotoxic T lymphocytes (CTL) obtained  
 30 from the HhD mice as described before (62).

Anti BA-46 peptide CTL activity showed variability in immunogenicity at a range between 20 % to 60 %. Lysis of 60 % was obtained against BA-46-7, 40 % against BA-46-6, BA-46-9 and 20 % against the rest of the peptides (Figure 14). CTL from HhD mice immunized with breast tumor extract peptide induced  
 35 10% to 30 % specific lysis of BA-46 peptide loaded RMA-S-HhD cells, 31 % lysis of tumor extract loaded targets and only 12 % and 14 % lysis of normal breast extracted peptides and colon extracted peptides loaded RMA-S-HhD cells, respectively. No lysis of the melanoma associated synthetic tyrosinase peptide

was obtained (Figure 15). The ability of BA-46 peptides to induce a breast associated CTL reaction that is HLA-A2.1 restricted was further examined. CTL against individual BA-46 peptides showed 30-50 % higher activity against breast tumor extract versus normal breast loaded target cells, supporting the fact that a preferential activity is obtained against breast tumor TAAs. These effectors also lysed preferentially a breast carcinoma line-HhD transfectant relative to parental cells, stressing the HLA-A2 restriction of the reaction (Figures 16 and 17).

#### EXAMPLE 4

##### 10      **Characterization of novel breast carcinoma MUC1 derived peptides as CTL epitopes**

The polymorphic epithelial Mucin, encoded by the MUC-1 gene is a high molecular weight transmembranal glycoprotein overexpressed in a broad range of tumors, such as breast, pancreas, ovary, thyroid and myeloma (72) cancer. It was found that the growth rate of primary breast tumors induced by the polyoma middle T antigen is significantly slower in MUC-1 null mice, suggesting that MUC-1 might play a role in the progression of mammary carcinoma (73). Furthermore, high level of MUC-1 expression by human breast cancer cells was found to be directly correlated with the presence of axially lymph node metastasis (74). A major feature of the MUC-1 molecule is the presence of an highly immunogenic extracellular tandem repeat array (TRA) heavily O-glycosidic-linked serine and threonine residues (75). Altered carbohydrate structure of MUC-1 in breast cancer cells, is probably responsible for the exposure of core epitopes within MUC-1, specifically recognized by monoclonal antibodies, as well as non MHC-restricted cytotoxic T lymphocytes (76). HLA-A11 (77) and more recently for HLA-A2.1 (78) restricted responses to the extracellular TRA have also been reported. However it was suggested that Mucin can actively suppress cell-mediated response against glycosylated TRA (79), or induced direct T cell apoptosis (80). More recently Agrawal et al showed that synthetic peptides derived from MUC-1 TRA cause suppression of human T-cell proliferative responses (81). This data points out the ambiguous role of MUC-1 TRA in T cell activation.

A series of peptides derived from non-TRA domains of the MUC-1 as potential CTL epitopes. The MUC-1 sequence was screened for potential HLA-A2.1 binding peptides (Table 8) and the eight peptides were synthesized.

TABLE 8

*Predicted human breast associated MUC-1 peptides that bind to HLA-A2*

Peptide	Sequence	Amino-acids*	SEQ ID:
MUC1-/C6	LLLLTVLTV	31-40	42
5 MUC1-/D6	LLLTVLTVV	32-41	43
MUC1-/F6	FLSFHISNL	323-331	44
MUC1-/A5	LLVLVCVLV	442-451	45
MUC1-/F4	ALLVLVCVL	441-450	46
MUC1-/B5	SLSYTNPAV	519-528	47
10 MUC1-/A7	NLTISDVSV	412-421	48
MUC1-/E6	ALASTAPPV	226-234	49

The prediction program is described in Materials and Methods. The peptides are listed in descending order of predicted HLA-A2.1-peptide complex stability. Single letter amino acid codes and the position of the first amino acid of the peptide in the protein sequence as well as the calculated binding score are listed. \*  
 15 Numbering is according to NCBI accession No. 182253.

These 9 residue peptides are derived from the signal peptide, cytoplasmic and extracellular domains of the MUC-1 protein. Among these peptides only the  
 20 MUC-1/B5 peptide has an identical sequence to the murine homologue. HLA-A.21 binding of MUC-1- derived peptides were evaluated by FACS analysis. The selected peptides were loaded on the murine TAP-deficient RMA-S-HhD transfectants and MHC stabilization was monitored (Figure 18). Although all peptides bound efficiently at the 1-100  $\mu$ M range, 3 peptides  
 25 MUC-1/D6, MUC-1/E6 and MUC-1/A7 exhibited higher binding affinity. In addition, similar binding affinities of these peptides were obtained upon loading on human TAP deficient T2 cells, expressing endogenous HLA-A2.1 molecules (data not shown).

Synthetic peptides corresponding to the MUC-1 TRA epitopes were shown  
 30 to induce CTL reaction in patients (77) as well as in HLA-A2.1K<sup>b</sup> transgenic mice (78). Hence we primarily examined the lysis pattern of each of the particular peptides listed in Table 8 following an immunization with a pool of the MUC-1-derived synthetic peptides (Figure 19). CTL results showed significant lysis of RMA-S-HhD target cells loaded either with the MUC-1/D6 peptide (38  
 35 %) or with the MUC-1/A7 peptide (30 %). Lower lysis rate of 15 % with the MUC-1/E6 and with the MUC-1/F6 peptides was demonstrated. The rest of the peptides showed only background lysis properties. Hence MUC-1/D6, MUC-1/A7 and MUC-1/E6 harbor immunogenic properties.

To determine the processing and presentation of MUC-1-derived peptides by breast carcinoma tumors, we selected the MDA-MB-157 cell line which is characterized by high MUC-1 expression and low class I expression (data not shown). Both the parental tumor cell line and its HhD transfectant (MDA-MB-157-HhD), were used as targets in CTL assays (Figure 20). Mice were immunized with either RMA-S-HhD-B7.1 cells loaded with MUC-1 selected peptides MUC-1/D6, MUC-1/A7 and MUC-1/E6, or with peptide extract derived from fresh patients' tumors. Preferential lysis of the MDA-MB-157-HhD cell line by anti MUC-1-derived peptides activated lymphocytes suggested both breast associated as well as MHC-restricted lysis. Moreover, this data strongly support the processing and presentation of non TRA associated MUC-1-derive peptides in MDA-MB-157-HhD breast carcinomas cells. Further analysis showed inhibition of lysis by anti HLA monoclonal antibody w6/32 (data not shown). In addition specific lysis of MDA-MB-157-HhD cells by CTL directed against fresh tumor extracted peptides emphasized an overlap in the peptide repertoire between the breast tumor cell line and fresh breast tumor extract.

To prove the existence of MUC-1/D6, MUC-1/A7 and MUC-1/E6 peptides in patients' derived breast tumor peptide extracts, a CTL experiment was performed by utilizing CTL against breast tumor peptide extracts as effectors against target cells presenting MUC-1 peptides or a control of normal breast tissue (Figure 21). All 3 MUC-1-derived peptides, but not the HLA-A2.1 melanoma associated peptide tyrosinase, could be recognized and lysed by anti-tumor extract CTL. RMA-S-HhD cells loaded with normal breast extract gave in different experiment between 40-50 % of the lysis induced against tumor extract loaded targets. This result is expected since a part of the tumor extract peptide repertoire consists of normal peptides.

A crucial parameter for selection of TAA peptides based vaccines are their expression frequency by tumors in comparison with normal tissues. Since MUC-1 protein is known to be overexpressed in tumors, with no tumor specific mutations, it is of obvious interest to examine the abundance of MUC-1 peptides in patients' derived normal breast tissue extract in comparison to tumor extract. CTL generated against MUC-1 peptides MUC-1/D6, MUC-1/A7 and MUC-1/E6, showed a 1.8-9.0 fold higher reactivity to tumor extract versus normal tissue extract (Figure 22). The same difference in preferential tumor versus normal tissue recognition by CTL could be detected upon vaccination with breast tumor extract peptides, supporting the window of specificity between normal and tumor tissues. These results suggest that MUC-1/D6, MUC-1/A7 and MUC-1/E6 are potential tumor associated antigen peptides.

## EXAMPLE 5

*Induction of CTL response by Teratocarcinoma-derived growth factor (CRIPTO-1) derived peptides.*

The CRIPTO-1 gene (also known as Teratocarcinoma-derived growth factor) was first identified and cloned from an undifferentiated human embryonal carcinoma cell line (83). It encodes a 188 amino acid glycoprotein with a region of structural homology to members of the epidermal growth factor family, but lacks a hydrophobic signal peptide and transmembrane domain. The receptor for CRIPTO-1 has not been identified, yet it has been shown to function as a growth factor. The gene can act as a dominantly transforming oncogene *in vitro*. A number of studies have shown elevated expression of CRIPTO-1 mRNA and protein in several human cancers, including human primary breast, gastric, colon, pancreatic, and bladder (84-88). It is therefore an attractive candidate tumor associated antigen for specific, active immunotherapy. Peptides have been synthesized with HLA-A2.01 binding potential based on the published sequences of CRIPTO-1, and these peptides have been used for immunization of HhD transgenic mice (see Table 9). Figures 24a-c show that some of the tested peptides (C1, C8, C10, and C12) can elicit lymphocyte response which specifically lyses CRIPTO-1 peptide-loaded target cells.

TABLE 9  
*Predicted human Cripto-1 derived peptides that bind to HLA-A2*

Peptide	Start Position	Sequence
Cripto-1/C1	5	KMARFSYSV
Cripto-1/C2	151	GLVMDEHLV
Cripto-1/C3	145	FLPGCDGLV
Cripto-1/C4	89	CMLGSFCAC
Cripto-1/C5	43	YLAFRDDSI
Cripto-1/C6	123	WLPKKCSLC
Cripto-1/C7	83	CLNGGTCML
Cripto-1/C8	176	MLVGICLSI
Cripto-1/C9	23	FELGLVAGL
Cripto-1/C10	5	KMVRFSYSV
Cripto-1/C11	83	CLNEGTCML
Cripto-1/C12	176	MLAGICLSI

Table 9 shows the sequences of the peptides tested in single letter amino acid code and their starting position in the intact protein (according to NCBI accession nos. 117473 (C1-C9) and 321120 (C10-C12)).

Although the invention has been described in conjunction with specific embodiments thereof, it is evident that many alternatives, modifications and variations will be apparent to those skilled in the art. Accordingly, it is intended



Variable	Mean	SD	Min	Max	Skewness	Kurtosis	Normality
Age	35.2	12.5	18	65	0.15	-0.5	0.98
Gender	1.2	0.4	1	2	0.05	-0.2	0.99
Education	12.5	2.1	9	16	0.25	-0.8	0.95
Income	45000	15000	20000	80000	0.35	-1.0	0.92
Health	2.5	0.8	1	4	0.10	-0.3	0.99
Stress	3.2	1.1	1	5	0.20	-0.7	0.96
Life Satisfaction	4.1	0.9	2	5	0.18	-0.6	0.97
Work Satisfaction	3.8	1.0	2	5	0.22	-0.8	0.94
Family Satisfaction	4.3	0.8	3	5	0.12	-0.4	0.98
Community Satisfaction	3.9	1.1	2	5	0.28	-0.9	0.91
Overall Satisfaction	4.0	0.9	2	5	0.16	-0.5	0.97

## REFERENCES CITED

1. Fidler, I.J. and Balch, C.M. (1987) The biology of cancer metastasis and implications for therapy. *Curr. Progr. Surg.* **24**, 137-209. Boring, C.C., Squires, T.S., Tong, T., Montgomery, S.,
2. Abeloff, M.D. (1996) Breast Cancer. *Current opinion in oncology* **8**, 447-448.
3. Andriole, G.L. (1997) Adjuvant therapy for prostate cancer patients at high risk of recurrence following radical prostatectomy. *Eur. Urol. Supp* **3**; 65-69.
4. Nseyo, U.O. and Lamm, D.L. (1997) Immunotherapy of bladder cancer. *Semin. Surg. Oncol.* **13**, 342-349.
5. Kedar, E. and Klein, E. (1995) Cancer Immunotherapy: Are the results discouraging? Can they be improved? *Adv. Immunol.* **58**, 245-322.
6. Urban, J.L. and Schreiber, H. (1992) Tumor antigens. *Annu. Rev. Immunol.* **10**, 617-644.
7. Van Pel, A. and Boon, T. (1982) Protection against a nonimmunogenic mouse leukemia by an immunogenic variant obtained by mutagenesis. *Proc. Natl. Acad. Sci. USA* **79**, 4718-4722.
8. Pardoll, D.M. (1993) New strategies for enhancing the immunogenicity of tumors. *Curr. Opinion Immunol.* **5**, 719-725.
9. Qin, Z. and Blankenstein, T. (1996) Influence of local cytokines on tumor metastasis: using cytokine gene transfected tumor cells as experimental models. In "Attempts to Understand Metastasis Formation III", *Therapeutic Approaches for Metastasis Treatment, Current Topics in Microbiology and Immunology* (U. Gunthert, P.M. Schlag and W. Birchmeier, eds.) **213III**, 55-64.
10. Pardoll, D.M. (1996) Cancer vaccines: a road map for the next decade. *Curr. Opinion Immunol.* **8**, 619-621.
11. Boon, T. and van der Bruggen, P. (1996) Human tumor antigens recognized by T lymphocytes. *J. Exptl. Med.* **183**, 725-729.
12. Townsend, A. and Bodmer, H. (1989) Antigen recognition by class I-restricted T lymphocytes. *Annu. Rev. Immunol.* **7**, 601-624.
13. Rammensee, H.-G., Falk, K. and Rotzschke, O. (1993) Peptides naturally presented by MHC class I molecules. *Annu. Rev. Immunol.* **11**, 213-244.
14. Feltkamp, M., Smits, H.L., Vierboom, M., Minnaar, R.P., deJough, B.M., Drijfhout, J.W., Schegget, J., Melief, C.J.M. and Kast, W.M. (1993) Vaccination with cytotoxic T lymphocyte epitope-containing peptide protects against a tumor induced by human papillomavirus type 16-transformed cells. *Eur. J. Immunol.* **23**, 2242-2249.

15. Kast, W.M., Brandt, R.M.P., Sidney, J., Drijfhout, J.W., Kubo, R.T., Melief, C.J.M. and Sette, A. (1994) Role of HLA-A motifs in identification of potential CTL epitopes in human papillomavirus type 16 E6 and E7 proteins. *J. Immunol.* **152**, 3904-3912.
16. Mandelboim, O., Berke, G., Fridkin, M., Feldman, M., Eisenstein, M. and Eisenbach, L. (1994) CTL induction by a tumor-associated antigen octapeptide derived from a murine lung carcinoma. *Nature (Lond.)* **369**, 67-71.
17. Cox, A., Skipper, J., Chen, Y., Henderson, R., Darrow, T., Shabanowitz, J., Engelhard, V., Hunt, D. and Slingluff, C. (1994) Identification of a peptide recognized by five melanoma-specific human cytotoxic T cell lines. *Science* **264**, 716-719.
18. Blake, J., Johnston, J., Hellstrom, K.E., Marquardt, H. and Chen, L. (1996) Use of combinatorial peptide libraries to construct functional mimics of tumor epitopes recognized by MHC class I restricted T lymphocytes. *J. exp. Med.* **184**, 121-130.
19. Boon, T., Cerottini, J.-C., Van den Eynde, B., Van der Bruggen, P. and Van Pel, A. (1994) Tumor antigens recognized by T lymphocytes. *Annu. Rev. Immunol.* **12**, 337-365.
20. Marchand, M., Weynants, P., Rankin, E., Arienti, F., Belli, F., Parmiani, G., Cascinelli, N., Bourland, A., Wanwijck, R., Humblet, Y., Canon, J.L., Naeyaert, J.M., Plagne, R., Deraemaeker, R., Knuth, A., Jager, E., Brasseur, F., Herman, J., Coulie, P.G. and Boon, T. (1995) *Int. J. Cancer* **63**, 883-885.
21. Rosenberg, S.A. et al. (1998) Immunologic and therapeutic evaluation of a synthetic peptide vaccine for the treatment of patients with metastatic melanoma. *Nature Med.* **4**, 321-327.
22. Nestle, F.L., Aljaghi, S., Gilliet, M., Sun, Y., Grabbe, S., Dummer, R., Burge, G. and Schadendorf, D. (1998) Vaccination of melanoma patients with peptide or tumor lysate-pulsed dendritic cells. *Nature Med.* **4**, 328-332.
23. Mandelboim, O., Vadai, E., Fridkin, M., Katz-Hillel, A., Feldman, M., Berke, G. and Eisenbach, L. (1995) Regression of established murine carcinoma metastases following vaccination with tumor-associated antigen. *Nature Med.* **1**, 1179-1183.
24. Mayordomo, J.I., Zorina, T., Storkus, W.J., Zitvogel, L., Cellzsi, C., Falo, L.D., Melief, C.J., Ildstad, W.M., Deleo, A.B. (1995) Bone marrow derived dendritic cells pulsed with synthetic tumor peptides elicit protective and therapeutic anti tumor immunity. *Nature Medicine* **1**, 1297-1302.

25. Fearon, E.R., Pardoll, D.M., Itaya, T., Golumbek, P., Levitsky, H.I., Simons, J.W., Karasuyama, H., Vogelstein, B., and Frost, P. (1990) Interleukin-2 production by tumor cells by passes T helper function in generation of an anti-tumor response. *Cell* **60**, 397-403.
26. Bakker, A., Schreurs, M., deBoer, A., Kawakami, Y., Rosenberg, S., Adema, G. and Figdor, C. (1994) Melanocyte lineage-specific antigen gp100 is recognized by melanoma-derived tumor-infiltrating lymphocytes. *J. Exp. Med.* **179**, 1005-1009.
27. Wentworth, P.A., Vitiello, A., Sidney, J., Keogh, E., Chesnut, R.W., Grey, H. and Sette, A. (1996) Differences and similarities in the A2.1-restricted cytotoxic T cell repertoire in humans and human leukocyte antigen-transgenic mice. *Eur. J. Immunol.* **26**, 97-101.
28. Barra, C., Gournier, H., Garcia, Z., Marche, P.N., Jouvin-Marche, E., Briand, P., Fillipi, P. and Lemonnier, F.A. (1993) Abrogation of H-2-restricted CTL responses and efficient recognition of HLA-A3 molecules in DBA/2 HLA/A24 responder mice. *J. Immunol.* **150**, 3681-3689.
29. Vitiello, A., Marchesini, D., Furze, J., Sherman, L.A., Chestnut, R.W. (1991) Analysis of HLA restricted influenza specific cytotoxic T lymphocytes response in transgenic mice carrying a chimeric human-mouse class I major histocompatibility complex. *J. Exp. Med.* **173**, 1007-1015.
30. Pascolo, S., Bervas, N., Ure, J.M., Smith, A.G., Lemmonier, F.A. and Rerarnau, B. (HLA-A2.1 restricted education and cytolytic activity of CD8<sup>+</sup> T lymphocytes from 2 microglobulin (2m) HLA-A2.1 monochain transgenic, H-2D<sup>b</sup>, 2m double knockout mice. *J. Exp. Med.* **185**, 2043-2051.
31. Spooner, R.A., Deonarain, M.P. and Epenetos, A.A., DNA vaccination for cancer treatment. *Gene Therapy* **2**, 173-180, 1995.
32. Ulmer, J.B., Donnelly, J.J., Parker, S.E., Rhodes, G.H., Felgner, P.L., Dwaraki, V.J., Gromkowski, S.H., Deck, R.R., DeWitt, C.M., Friedman, A., Hawe, L.A., Leander, K.R., Martinez, D., Perry, H.C., Shiver, J.W., Montgomery, D.L. and Liu, M.A., Heterologous protection against influenza by injection of DNA encoding a viral protein. *Science* **259**, 1745-1749, 1993.
33. Conry, R.M., LoBuglio, A.F., Kantor, J., Schlom, J., Loechel, F., Moore, S.E., Sumerel, L.A., Barlow, D.L., Abrams, S. and Curiel, D.T., Immune response to a carcinoembryonic antigen polynucleotide vaccine. *Cancer Res.*, **54**, 1164-1168, 1994.

34. Conry, R.M., LoBuglio, A.F., Loechel, F., Moore, S.E., Sumerel, L.A., Barlow, D.L., Pike, J. and Curiel, D.T. A carcinoembryonic antigen polynucleotide vaccine for human clinical use. *Cancer Gene Therapy* 2, 33-38, 1995.
35. Acres, R.B., Hareuveni, M., Balloul, J.-M. and Kieny, M.-P. Vaccinia virus MUC1 immunization of mice: immune response and protection against the growth of murine tumors bearing the MUC1 antigen. *J. Immunother.* 14, 136-143, 1993.
36. Whitton, J.L., Sheng, N., Oldstone, M.B. and McKee, T.A., A "string-of-beads" vaccine, comprising linked minigenes, confers protection from lethal-dose virus challenge. *J. Virol.* 67, 348-352, 1993.
37. Thomson, S.A., Elliott, S.L., Sherritt, M.A., Sproat, K.W., Coupar, B.E., Scalzo, A.A., Forbes, C.A., Ladhams, A.M., Mo, X.Y., Tripp, R.A., Doherty, P.C., Moss, D.J. and Suhrbier, A., Recombinant polypeptide vaccines for the delivery of multiple CD8 cytotoxic T cell epitopes. *J. Immunol.* 157, 822-826, 1996.
38. Thomson, S.A., Sherritt, M.A., Medveczky, J., Elliott, S.L., Moss, D.J., Fernando, G.J., Brown, L.E., and Suhrbier, A., Delivery of multiple CD8 cytotoxic T cell epitopes by DNA vaccination. *J. Immunol.* 160, 1717-1723, 1998.
39. Boczkowski, D., Nair, S.K., Snyder, D. and Gilboa, E., Dendritic cells pulsed with RNA are potent antigen-presenting cells in vitro and in vivo. *J. Exp. Med.* 184, 465-472, 1996.
40. Mayordomo, J.I., Zorina, T., Storkus, W.J., Zitvogel, L., Celluzzi, C., Falo, L.D., Melief, C.J., Ildstad, S.T., Kast, W.M., Deleo, A.B., et al., Bone marrow-derived dendritic cells pulsed with synthetic tumor peptides elicit protective and therapeutic antitumor immunity. *Nature Med.* 1, 1297-1302, 1995.
41. Mandelboim, O., Berke, G., Fridkin, M., Feldman, M., Eisenstein, M., and Eisenbach, L., CTL induction by a tumor-associated antigen octapeptide derived from a murine lung carcinoma. *Nature* 369, 67-71, 1994.
42. Mandelboim, O., Vadai, E., Fridkin, M., Katz-Hillel, A., Feldman, M., Berke, G., and Eisenbach, L., Regression of established murine carcinoma metastases following vaccination with tumour-associated antigen peptides. *Nat. Med.* 1, 1179-1183, 1995.
43. Mukherji, B., Chakraborty, N.G., Yamasaki, S., Okino, T., Yamase, H., Sporn, J.R., Kurtzman, S.K., Ergin, M.T., Ozols, J., Meehan, J., et al., Induction of antigen-specific cytolytic T cells in situ in human

- melanoma by immunization with synthetic peptide-pulsed autologous antigen presenting cells. *Proc. Natl. Acad. Sci. USA* 92, 8078-8082, 1995.
44. Gjertsen, M.K., Bakka, A., Breivik, J., Saeterdal, I., Gedde-Dahl, Stokke, K.T., Solheim, B.G., Egge, T.S., Soreide, O., Thorsby, E., and Gaudernack, G., Ex vivo ras peptide vaccination in patients with advanced pancreatic cancer: results of a phase I/II study. *Int. J. Cancer* 65, 450-453, 1996.
  45. Murphy, G., Tjoa, B., Ragde, H., Kenny, G. and Boynton, A., Phase I clinical trial: T-cell therapy for prostate cancer using autologous dendritic cells pulsed with HLA-A0201-specific peptide from prostate-specific membrane antigen. *Prostate* 29, 371-380, 1996.
  46. Messing, E.M. and Catalona, W. Urothelial tumors of the urinary tract. In *Campbell's Urology*, Eds. Walsh, P.C., Retik, A.B., Vaughan, E.D. and Wein, A.J. (7th editin, WB, Saunders Co. 1998), 2327-2410.
  47. Ioachim-Velogianni, E., Stavropoulos, N.E., Kitsiou, E., Stefanaki, S. and Agnantis, N.J. (1995) Distribution of CD1A-positive Langerhans cells and lymphocytes subsets in transitional cell carcinoma of the urinary bladder. An immunohistological study on frozen sections. *J. Pathol.* 155, 401-406.
  48. Thanhauser, A., Bohle, A., Flad, H.D., Ernst, M., Mattern, T. and Ulmer, A.J. (1993) Induction of bacillus Calmette-Gherin activated killer cells from human peripheral blood mononuclear cells against bladder carcinoma cell lines *in vitro*. *Cancer Immunol. Immunother.* 37, 105-111.
  49. Patard, H., Muscatelli-Groux, B., Saint, F., Popov, Z., Maille, P., Abbou, C. and Chopin, D. (1996) Evaluation of local immune response after intravesical bacille Calmette Guerin treatment for superficial bladder cancer. *Br. J. Urol.* 78, 709-714.
  50. Housseau, F., Zeliskewski, D., Roy, M. et al. (1997) MHC-dependent cytotoxicity of autologous tumor cells by lymphocytes infiltrating urothelial carcinomas. *Int. J. Cancer* 71, 585-594.
  51. Moll, R., Wu, X-R., Lin, J-H. and Sun, T-T. (1995) Uroplakins, specific membrane proteins of urothelial umbrella cells, as histological markers of metastatic transitional cell carcinomas. *Am. J. Pathol.* 147, 1383-1397.
  52. Mandelboim, O., Bar-Haim, E., Vadai, E., Fridkin, M. and Eisenbach, L. (1997) Identification of shared Tumor-Associated Antigen peptides between two spontaneous lung carcinomas. *J. Immunol.* 159, 6030-6036.

53. Boring, C.C., Squires, T.S., Tong, T. and Montgomery, S. (1994) Cancer statistics 1994 CA 43, 7-26.
54. Carter, H.B., Pearson, J.D., Metter, J.E. et al. (1992) Longitudinal evaluation of prostate specific antigen levels in men with and without prostate disease. JAMA 267, 2215-2220.
55. Partin, A.W., Poind, C.R., Clemens, J.Q. et al. (1993) Serum PSA after anatomic radical prostatectomy. Urol. Clin. North Am. 20, 713-725.
56. Epstein, J.I., Partin, A.W., Sauvageot, J. and Walsh, P.C. (1996) Prediction of progression following radical prostatectomy. A multivariate analysis of 721 men with long term follow up. Am. J. Surg. Pathol. 20, 286-292.
57. Eisenberger, M.H. (1998) Chemotherapy for hormone resistant prostate cancer. In Campbell's Urology, Eds. Walsh, P.C., Retik, A.B., Vaughan, E.D. and Wein, A.J. (7th edition, WB, Saunders Co.) 2645-2658.
58. Schroder, F.H. (1998) Endocrine treatment of prostate cancer. In Campbell's Urology, Eds. Walsh, P.C., Retik, A.B., Vaughan, E.D. and Wein, A.J. (7th edition, WB, Saunders Co.) 2627-2644.
59. Xue, B.H., Zhang, Y., Sosman, J.A. and Peace, D.J. (1997) Induction of human cytotoxic T lymphocytes specific for prostate specific antigen. The Prostate 30, 73-78.
60. Correale, P., Walmsley, K., Nieroda, C., Zhu, M., Schlom, J. and Tsang, K.Y. (1997) *In vitro* generation of human cytotoxic T lymphocytes specific for peptides derived from prostate specific antigen. J. Natl. Cancer Inst. 89, 272-275.
61. Tjoa, B., Bounton, A., Kenny, G., Ragde, H., Misrock, S.L. and Murphy, G. (1996) Presentation of prostate tumor antigens by dendritic cells stimulates T cell proliferation and cytotoxicity. Prostate 28, 65-69.
62. Murphy, G., Tjoa, B., Ragde, H., Kenny, G. and Bounton, A. (1996) Phase I clinical trial: T cell therapy for prostate cancer using autologous dendritic cells pulsed with HLA-A0201 specific peptides from Prostate Specific Membrane Antigen. The Prostate 29, 371-380.
63. Bonandon, G. (1996) Future development in adjuvant systemic therapy for high risk breast cancer. Cancer Res. 140, 227-234.
64. Barnd, D.L., Lan, M., Metzger, R., Finn, O.J. (1989). Specific, MHC-unrestricted recognition of tumor associated mucins by human cytotoxic T cells. Proc. Natl. Acad. SCI. USA 86, 7159-7163.
65. Agrawal, B., Reddish, M., Longenecker, B.M. (1996). *In vitro* induction of MUC-1 peptide specific type 1 T lymphocyte and cytotoxic T lymphocyte responses from healthy multiparous donors. J. Immunol 157, 2089-2095.

66. Fisk, B., Blevins, T.L., Taylor-Wharton, J., Ionnides, C.G., (1995). Identification of an immunodominant peptide of HER2/neu protooncogene recognized by ovarian Tumor specific cytotoxic T lymphocyte lines. *J. Exp. Med.* **181**, 2109-2117.
67. Linehan, C., Goedegebuure, P.S., peoples, G.E., Rogers, S.O., Eberlein, P.J. (1995) Tumor specific and HLA-A2 restricted cytotoxicity by tumor-associated lymphocytes in human metastatic breast cancer. *J. Immunol* **155**, 4486-4491.
68. Larocca, D., Peterson, J.A., Urrea, R., Kuniyoshi, J., Bistrain, A.M. and Ceriani, R.L. (1991) A Mr 46,000 human milk fat globule protein that is highly expressed in human breast tumors contains factor VIII-like domains. *Cancer Res.* **51**, 4994-4998.
69. Taylor, M.R., Couto, J.R., Scallan, C.D., Ceriani, R.L. and Peterson, J.A. (1997) Lactadherin (formerly BA-46), a membrane-associated glycoprotein expressed in human milk and breast carcinomas, promotes Arg-Gly-Asp (RGD)-dependent cell adhesion. *DNA Cell Biol.* **16**, 861-869.
70. Ceriani, R.L., Blank, E.W., Couto, J.R., and Peterson, J.A. (1995) Biological activity of two humanized antibodies against two different breast cancer antigens and comparison to their original murine forms. *Cancer Res.* **55**, 5852-5856S.
71. Couto, J.R., Blank, E.W., Peterson, J.A., Ceriani, and R.L. (1995) Anti-BA-46 monoclonal antibody Mc3: humanization using a novel positional consensus and *in vivo* and *in vitro* characterization. *Cancer Res.* **55**, 1717-1722.
72. Ho, S.B., Niehans, G.A., Lyftogt, C., Yan, P.S., Cherwitz, D.L., Gum, E.T., Dahiya, R., and Kim, Y.S. (1993) Heterogeneity of Mucin gene expression in normal and neoplastic tissues. *Cancer Res* **53**, 641-651.
73. Spicer, A.P., Rowse, G.J., Linder, T.K. and Gendler, S.J. (1995) Delayed mammary tumor progression in Muc-1 null mice. *J. Biol. Chem.* **270**, 30093.
74. McGuckin, M.A., Walsh, M.D., Hohn, B.G., Ward, B.G. and Wright, R.G. (1995) Prognostic significance of Muc-1 epithelial Mucin expression in breast cancer. *Hum. Pathol.* **26**, 432.
75. Wreschner, D.H., Hareuveni, M., Tsarfaty, I., Smorodinsky, N., Horev, J., Zaretsky, J., Kotkes, P., Weiss, M., Lathe, R. and Dion, A. (1990) Human epithelial tumor antigen sequences. Differential splicing may generate multiple protein forms. *Eur. J. Biochem.* **189**, 463.
76. Domenech, N. and Finn, O.J. (1995) *In vitro* studies of MHC-restricted recognition of a peptide from the MUC1 tandem repeat domain by CTL. *FASEB J.* **9**, 1023.



77. Domenech, N., Henderson, R.A. and Finn, O.J. (1995) Identification of a HLA-A11-restricted epitope from the tandem repeat domain of the epithelial tumor antigen Mucin. *J Immunol* **15**, 4766.
78. Apostolopoulos, V., Karanikas, V., Haurum, J.S. and McKenzie, I.F. (1997) Induction of HLA-A2-restricted CTLs to the Mucin 1 human breast cancer antigen. *J. Immunol.* **159**, 5211.
79. Mackenzi, P.Y. and Longenecker, B.M. (1991) Specific immunosuppressive activity of epiglycanin, a Mucin-like glycoprotein secreted by a murine mammary adenocarcinoma (TA3-HA). *Cancer Res.* **15**, 1170.
80. Gimmi, C.D., Morrison, B.W., Mainpric, B.A., Gribben, J.G., Boussiotis, V.A., Freeman, G.J., Park, S.Y., Watanabe, M., Gong, J., Hayes, D.F., Kufe, D.W. and Nadler, L.M. (1996) Breast cancer-associated antigen, DF3/MUC1, induces apoptosis of activated human T cells. *Nat. Med.* Dec. **2**, 1367.
81. Agrawal, B., Krantz, M.J., Reddish, M.A., Longenecker and B.M. (1998) Cancer-associated MUC1 Mucin inhibits human T-cell proliferation, which is reversible by IL-2. *Nat Med.* **4**, 43.
82. Parker, K.C., Bednarek, M.A., Coligan, J.E., Scheme for ranking potential HLA-A2 binding peptides based on independent binding of individual peptide side-chains. *J Immunol.* **152**, 163-175, 1994.
83. Ciccodicola, A., Dono, R., Obici, S., Simeone, A., Zollo, M., and Persico, M.G. (1989) expressed in undifferentiated human NTERA2 teratocarcinoma cells. *EMBO J* **8**, 1987.
84. Qi, C. F., Liscia, D. S., Normano, N., Merlo, G., Johnson, G.R., Gullick, W.J., Ciardiello, F., Saeki, T., Brandt, R., Kim, N., Kenney, N., and Salomon, D.S.. (1994) Expression of Transforming Growth Factor A, Amphiregulin and Cripto-I in Human Breast Carcinomas. *Br J Cancer*, **69**, 903.
85. Kuniyasu, H., Yoshida, K., Yokoazaki, H., et al. (1991) Expression of Cripto, a Novel Gene of the Epidermal Growth Factor Family, in Human Gastrointestinal Carcinomas. *Jpn J Cancer Res* **82**, 969.
86. Saeki, T., Stromberg, K., Qi, C.F., et al. (1992) Differential Immunohistochemical Detection of Amphiregulin and Cripto in Human Normal Colon and Colorectal Cancers Tumours. *Cancer Res.* **52**, 3467.
87. Freiss, H., Yamanaka, Y., Buchler, M., Korbin, M. S., et al. (1994) Cripto, A Member of The Epidermal Growth Factor Family, is Over Expressed in Human Pancreatic Cancer and Chronic Pancreatitis. *Int J Cancer* **56**, 688.

88. Byrne, R.L., Autzen, P., Birch, P., Robinson, M.C., Gullick, W.J., et al. (1998). The Immunohistochemical Detection of Cripto-1 in Benign and Malignant Human Bladder. *J. Pathol.* **185**, 108.

ART 34 AMDT

54

## WHAT IS CLAIMED IS

1. A peptide derived from a protein selected from the group consisting of Uroplakin (UP), Prostate specific antigen (PSA), Prostate specific membrane antigen (PSMA), Prostate acid phosphatase (PAP), Lactadherin (BA-46), Mucin (MUC1) and Teratocarcinoma-derived growth factor (CRIPTO-1), the peptide comprising 8 to 10 amino acid residues, of which a second residue from an amino terminal of the peptide and an end residue at a carboxy terminal of the peptide are hydrophobic or hydrophilic natural or non-natural amino acid residues, with the proviso that for PSA, SEQ ID Nos 20 and 24 are excluded, for PSMA, SEQ ID Nos 25, 26, 27, 29 and 30 are excluded and for PAP, SEQ ID Nos 31, 32, 33 and 34 are excluded.
2. The peptide of claim 1, wherein the peptide is derived from Uroplakin.
3. The peptide of claim 2, wherein said Uroplakin is selected from the group consisting of Uroplakin II, Uroplakin Ia, Uroplakin III and Uroplakin Ib.
4. The peptide of claim 3, wherein the peptide has a sequence selected from the group consisting of SEQ ID NOs:1-19 and 50-64.
5. The peptide of claim 1, wherein the peptide is derived from Prostate specific antigen (PSA) but excluding SEQ ID Nos 20 and 24.
6. The peptide of claim 5, wherein the peptide has a sequence selected from the group consisting of SEQ ID NO's 21-23.
7. The peptide of claim 1, wherein the peptide is derived from Prostate specific membrane antigen (PSMA) but excluding SEQ ID Nos 25, 26, 27, 29 and 30.
8. The peptide of claim 7, wherein the peptide has the sequence identified as SEQ ID NO 28.

9. The peptide of claim 1, wherein the peptide is derived from Prostate acid phosphatase (PAP) but excluding SEQ ID Nos 31, 32, 33 and 34.
10. The peptide of claim 1, wherein the peptide is derived from said Mucin.
11. The peptide of claim 10, wherein the peptide is derived from a non-tandem repeat array of said Mucin.
12. The peptide of claim 10, wherein the peptide is derived from a region selected from the group consisting of a signal peptide, a cytoplasmic domain and an extracellular domain of said Mucin.
13. The peptide of claim 12, wherein the peptide is derived from a non tandem repeat array of said Mucin.
14. The peptide of claim 10, wherein the peptide has a sequence selected from the group consisting of SEQ ID NOs:42-49.
15. The peptide of claim 1, wherein the peptide is derived from said Lactadherin (BA-46).
16. The peptide of claim 15, wherein the peptide has a sequence selected from the group consisting of SEQ ID NOs:35-41.
17. The peptide of claim 1, wherein the peptide is derived from said Teratocarcinoma-derived growth factor (CRIPTO-1).
18. The peptide of claim 17, wherein the peptide has the sequence selected from the group consisting of SEQ ID Nos. 66-77.
19. The peptide of any of claims 1-18, wherein said peptide is derived from a mammal.
20. The peptide of claim 19, wherein the mammal is a humanoid or a rodent.

21. The peptide of any of claims 1-20, wherein said peptide includes at least one non-natural modification.
- 5 22. The peptide of claim 21, wherein said non-natural modification renders the peptide more immunogenic or more stable than the unmodified peptide.
- 10 23. The peptide of claim 21 or 22, wherein said at least one modification is selected from the group consisting of peptoid modification, semipeptoid modification, cyclic peptide modification, N terminus modification, C terminus modification, peptide bond modification, backbone modification and residue modification.
- 15 24. A pharmaceutical composition comprising, as an active ingredient, at least one peptide as set forth in any of claims 1-23 and a pharmaceutically acceptable carrier.
- 20 25. The pharmaceutical composition of claim 24, wherein said carrier is selected from the group consisting of a proteinaceous carrier to which said at least one tumor associated antigen peptide is linked, an adjuvant, a protein or a recombinant protein and an antigen presenting cell.
- 25 26. The pharmaceutical composition of claim 24, wherein the composition contains an amount of said peptide effective to prevent or cure cancer or cancer metastases.
- 30 27. The pharmaceutical composition of claim 26, wherein said cancer is selected from the group consisting of breast, bladder, prostate, pancreas, ovary, thyroid, colon, stomach and head and neck cancer.
28. The pharmaceutical composition of claim 26, wherein said cancer is a carcinoma.
29. The pharmaceutical composition of claim 24, wherein the composition is a vaccine.

30. A vaccine composition comprising, as an active ingredient, at least one peptide as set forth in any of claims 1-23 and a suitable carrier.
- 5 31. The vaccine composition of claim 30, wherein said carrier is selected from the group consisting of a proteinaceous carrier to which said at least one tumor associated antigen peptide is linked, an adjuvant, a protein or a recombinant protein and an antigen presenting cell.
- 10 32. The vaccine composition of claim 30, wherein the composition contains an amount of said peptide effective to prevent or cure cancer or cancer metastases.
- 15 33. The vaccine composition of claim 32, wherein said cancer is selected from the group consisting of breast, bladder, prostate, pancreas, ovary, thyroid, colon, stomach and head and neck cancer.
- 20 34. The vaccine composition of claim 32, wherein said cancer is a carcinoma.
- 25 35. A method of prevention or cure of a cancer or of metastases thereof comprising the step of administering to a patient an effective amount of the pharmaceutical composition of any of claims 24-29.
- 30 36. A method of prevention or cure of a cancer or of metastases thereof comprising the step of vaccinating a patient with an effective amount of the vaccine composition of any of claims 30-34.
37. A polynucleotide encoding at least one peptide according to any of claims 1-23.
38. A polynucleotide encoding at least one peptide selected from the group consisting of SEQ ID Nos. 1-19, 21-23, 28, 35-64 and 66-77.
39. The polynucleotide of claim 37 or 38, wherein the polynucleotide forms a part of a longer polynucleotide designed to encode a fused protein product from which said at least one peptide is cleavable by a protease.

40. A pharmaceutical composition comprising, as an active ingredient, at least one polynucleotide as set forth in any claims 37-39 and a pharmaceutically acceptable carrier.

41. A cellular vaccine composition comprising an antigen presenting cell presenting at least one peptide of any of claims 1-23.

42. The cellular vaccine composition of claim 41, wherein said antigen presenting cell is selected from the group consisting of a dendritic cell, a macrophage, a B cell and a fibroblast.

43. The cellular vaccine composition of claim 41, wherein said antigen presenting cell is caused to present said at least one tumor associated antigen peptide by a method selected from the group consisting of:

- a) genetically modifying said antigen presenting cell with at least one polynucleotide encoding said at least one tumor associated antigen peptide such that said peptide or at least one longer polypeptide including said peptide will be expressed;
- b) loading said antigen presenting cell with at least one polynucleotide encoding said at least one tumor associated antigen peptide;
- c) loading said antigen presenting cell with said at least one tumor associated antigen peptide; and
- d) loading said antigen presenting cell with at least one longer polypeptide including said at least one tumor associated antigen peptide

44. The peptide of claim 1, wherein the second residue and the end residue are neutral, hydrophobic and aliphatic.

45. The pharmaceutical composition of any of claims 24-29 or 40 also comprising a helper peptide.

46. The pharmaceutical composition of claim 45, wherein the helper peptide has a T helper epitope.


47. The vaccine composition of any of claims 30-34 also comprising a helper peptide.
48. The vaccine composition of claim 47, wherein the helper peptide has a T helper epitope.
49. Use of at least one peptide of claims 1-23 in the manufacture of a medicament.
50. The at least one peptide of claims 1-23 for use as a medicament.
51. The use of claim 49 or 50, wherein the medicament is effective to prevent or cure a cancer or cancer metastases.
52. A peptide derived from a protein selected from the group consisting of Uroplakin (UP), Lactadherin (BA-46), Prostate specific antigen (PSA), Prostate specific membrane antigen (PSMA), Prostate acid phosphatase (PAP), Mucin (MUC1) and Teratocarcinoma-derived growth factor (CRIPTO-1), the peptide comprising 8-10 amino acid residues selected so as to promote effective binding to a MHC class 1 type molecule so as to elicit a CTL response.

*PCamywp/048*




1/29

Fig. 1




2/29

Fig. 2




3/29


Fig. 3



4/29

Fig. 4





6/29

Fig. 6






Fig. 7




8/29

Fig. 8






9/29



10/29

Fig. 10



11/29




Fig. 11

408474804

-- 12/29




Fig. 13




Fig. 14




Fig. 15





Fig. 16





17/29




Fig.18




Fig. 19





Fig. 20






Fig. 21




Fig. 22

Fig. 23a






Fig. 23b



24/29


Fig. 23c






25/29

Fig. 23d




26/29

Fig. 23e




27/29

Fig. 24a




28/29

Fig. 24b



29/29

Fig. 24c



# Combined Declaration for Patent Application and Power of Attorney

As a below-named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name; and that I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

TUMOR ASSOCIATED ANTIGEN PEPTIDES AND USE OF SAME AS ANTI-TUMOR VACCINES

the specification of which (check one)

- [ ] is attached hereto;  
 [ ] was filed in the United States under 35 U.S.C. §111 on \_\_\_\_\_, as  
 U.S. Appln. No. \_\_\_\_\_\*; or  
 [X] was/will be filed in the U.S. under 35 U.S.C. §371 by entry into the U.S. national stage of an  
 international (PCT) application, PCT/IL99/00417, filed 29 July 1999, entry requested on  
 \_\_\_\_\_\*; national stage application received U.S. Appln. No. \_\_\_\_\_\*, §371/§102(e)  
 date \_\_\_\_\_\* (\* if known)

and was amended on \_\_\_\_\_ (if applicable).

(include dates of amendments under PCT Art. 19 and 34 if PCT)

I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above; and I acknowledge the duty to disclose to the Patent and Trademark Office (PTO) all information known by me to be material to patentability as defined in 37 C.F.R. §1.56.

I hereby claim foreign priority benefits under 35 U.S.C. §§ 119 and 365 of any prior foreign application(s) for patent or inventor's certificate, or prior PCT application(s) designating a country other than the U.S., listed below with the "Yes" box checked and have also identified below any such application having a filing date before that of the application on which priority is claimed:

<u>125608</u>	<u>Israel</u>	<u>30 July 1998</u>	[X]	[ ]
(Number)	(Country)	(Day Month Year Filed)	YES	NO
_____	_____	_____	[ ]	[ ]
(Number)	(Country)	(Day Month Year Filed)	YES	NO

I hereby claim the benefit under 35 U.S.C. §120 of any prior U.S. non-provisional application(s) or prior PCT application(s) designating the U.S. listed below, or under §119(e) of any prior U.S. provisional applications listed below, and, insofar as the subject matter of each of the claims of this application is not disclosed in such U.S. or PCT application in the manner provided by the first paragraph of 35 U.S.C. §112, I acknowledge the duty to disclose to the PTO all information as defined in 37 C.F.R. §1.56(a) which occurred between the filing date of the prior application and the national filing date of this application:

_____	_____	_____
(Application No.)	(Day Month Year Filed)	(Status: patented, pending, abandoned)
_____	_____	_____
(Application No.)	(Day Month Year Filed)	(Status: patented, pending, abandoned)
_____	_____	_____
(Application No.)	(Day Month Year Filed)	(Status: patented, pending, abandoned)

As a named inventor, I hereby appoint the following registered practitioners to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith:

**All of the practitioners associated with Customer Number 001444**

Direct all correspondence to the address associated with **Customer Number 001444**, i.e.

**BROWDY AND NEIMARK, P.L.L.C.**  
**624 Ninth Street, N.W.**  
**Washington, D.C. 20001-5303**  
**(202) 628-5197**

The undersigned hereby authorizes the U.S. Attorneys or Agents appointed herein to accept and follow instructions from Yeda Research and Development Co. Ltd. as to any action to be taken in the U.S. Patent and Trademark Office regarding this application without direct communication between the U.S. Attorneys or Agents and the undersigned. In the event of a change of the persons from whom instructions may be taken, the U.S. Attorneys or Agents appointed herein will be so notified by the undersigned.

Title: TUMOR ASSOCIATED ANTIGEN PEPTIDES AND USE OF SAME AS ANTI-TUMOR VACCINES

U.S. Application filed \_\_\_\_\_, Serial No. \_\_\_\_\_

PCT Application filed 29 July 1999, Serial No. PCT/IL99/00417

I hereby further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C. §1001 and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

FULL NAME OF FIRST INVENTOR <u>Lea EISENBACH</u>		INVENTOR'S SIGNATURE <u>Lea Eisenbach</u>	DATE <u>16/01/2001</u>
RESIDENT <u>Rehovot, Israel</u>		CITIZENSHIP <u>Israeli</u> <u>ILX</u>	
POST OFFICE ADDRESS <u>Hanassi Harishon Street 33, 76303 Rehovot, Israel</u>			
FULL NAME OF SECOND JOINT INVENTOR <u>Lior CARMON</u>		INVENTOR'S SIGNATURE	DATE
RESIDENT <u>Tel-Aviv, Israel</u>		CITIZENSHIP <u>Israeli</u>	
POST OFFICE ADDRESS <u>Kehilat-Kiyov Street 6, 69989 Tel-Aviv, Israel</u>			
FULL NAME OF THIRD JOINT INVENTOR <u>Boaz TIROSH</u>		INVENTOR'S SIGNATURE <u>Boaz Tirosh</u>	DATE <u>16/01/01</u>
RESIDENT <u>Kiryat Ono, Israel</u>		CITIZENSHIP <u>Israeli</u> <u>ILX</u>	
POST OFFICE ADDRESS <u>Stern Street 9, 55602 Kiryat Ono, Israel</u>			
FULL NAME OF FOURTH JOINT INVENTOR <u>Erez BAR-HAIM</u>		INVENTOR'S SIGNATURE <u>Erez Bar-Haim</u>	DATE <u>16/01/01</u>
RESIDENT <u>Yavne, Israel</u>		CITIZENSHIP <u>Israeli</u> <u>ILX</u>	
POST OFFICE ADDRESS <u>Hatsivony Street 25/8, 81573 Yavne, Israel</u>			
FULL NAME OF FIFTH JOINT INVENTOR <u>Adrian PAZ</u>		INVENTOR'S SIGNATURE	DATE
RESIDENT <u>Petach Tikva, Israel</u>		CITIZENSHIP <u>Israeli</u>	
POST OFFICE ADDRESS <u>Wolfson Street 25, 49541 Petach Tikva, Israel</u>			
FULL NAME OF SIXTH JOINT INVENTOR <u>Matityahu FRIDKIN</u>		INVENTOR'S SIGNATURE <u>Matityahu Fridkin</u>	DATE <u>18/01/2001</u>
RESIDENT <u>Rehovot, Israel</u>		CITIZENSHIP <u>Israeli</u> <u>ILX</u>	
POST OFFICE ADDRESS <u>Miller Street 23, 76284 Rehovot, Israel</u>			
FULL NAME OF SEVENTH JOINT INVENTOR <u>Cheryl FITZER-ATTAS</u>		INVENTOR'S SIGNATURE	DATE
RESIDENT <u>Rehovot, Israel</u>		CITIZENSHIP <u>US/Israeli</u>	
POST OFFICE ADDRESS <u>Sapir Street 9, 76227 Rehovot, Israel</u>			

#3

## Combined Declaration for Patent Application and Power of Attorney

As a below-named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name; and that I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

TUMOR ASSOCIATED ANTIGEN PEPTIDES AND USE OF SAME AS ANTI-TUMOR VACCINES

the specification of which (check one)

- [ ] is attached hereto;  
[ ] was filed in the United States under 35 U.S.C. §111 on \_\_\_\_\_, as  
U.S. Appln. No. \_\_\_\_\_\*; or  
[X] was/will be filed in the U.S. under 35 U.S.C. §371 by entry into the U.S. national stage of an  
international (PCT) application, PCT/IL99/00417; filed 29 July 1999, entry requested on  
\_\_\_\_\_\*; national stage application received U.S. Appln. No. \_\_\_\_\_\*; §371/§102(e)  
date \_\_\_\_\_\* (\* if known)

and was amended on \_\_\_\_\_ (if applicable).

*(include dates of amendments under PCT Art. 19 and 34 if PCT)*

I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above; and I acknowledge the duty to disclose to the Patent and Trademark Office (PTO) all information known by me to be material to patentability as defined in 37 C.F.R. §1.56.

I hereby claim foreign priority benefits under 35 U.S.C. §§ 119 and 365 of any prior foreign application(s) for patent or inventor's certificate, or prior PCT application(s) designating a country other than the U.S., listed below with the "Yes" box checked and have also identified below any such application having a filing date before that of the application on which priority is claimed:

<u>125608</u>	<u>Israel</u>	<u>30 July 1998</u>	[X]	[ ]
(Number)	(Country)	(Day Month Year Filed)	YES	NO
_____	_____	_____	[ ]	[ ]
(Number)	(Country)	(Day Month Year Filed)	YES	NO

I hereby claim the benefit under 35 U.S.C. §120 of any prior U.S. non-provisional application(s) or prior PCT application(s) designating the U.S. listed below, or under §119(e) of any prior U.S. provisional applications listed below, and, insofar as the subject matter of each of the claims of this application is not disclosed in such U.S. or PCT application in the manner provided by the first paragraph of 35 U.S.C. §112, I acknowledge the duty to disclose to the PTO all information as defined in 37 C.F.R. §1.56(a) which occurred between the filing date of the prior application and the national filing date of this application:

_____ (Application No.)	_____ (Day Month Year Filed)	_____ (Status: patented, pending, abandoned)
_____ (Application No.)	_____ (Day Month Year Filed)	_____ (Status: patented, pending, abandoned)
_____ (Application No.)	_____ (Day Month Year Filed)	_____ (Status: patented, pending, abandoned)

As a named inventor, I hereby appoint the following registered practitioners to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith:

**All of the practitioners associated with Customer Number 001444**

Direct all correspondence to the address associated with **Customer Number 001444**; i.e.,

**BROWDY AND NEIMARK, P.L.L.C.**  
**624 Ninth Street, N.W.**  
**Washington, D.C. 20001-5303**  
**(202) 628-5197**

The undersigned hereby authorizes the U.S. Attorneys or Agents appointed herein to accept and follow instructions from Yeda Research and Development Co. Ltd. as to any action to be taken in the U.S. Patent and Trademark Office regarding this application without direct communication between the U.S. Attorneys or Agents and the undersigned. In the event of a change of the persons from whom instructions may be taken, the U.S. Attorneys or Agents appointed herein will be so notified by the undersigned.



Title: TUMOR ASSOCIATED ANTIGEN PEPTIDES AND USE OF SAME AS ANTI-TUMOR VACCINES

U.S. Application filed \_\_\_\_\_, Serial No. \_\_\_\_\_

PCT Application filed 29 July 1999, Serial No. PCT/IL99/00417

I hereby further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C. §1001 and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

FULL NAME OF FIRST INVENTOR <b>Lea EISENBACH</b>	INVENTOR'S SIGNATURE	DATE
RESIDENT <b>Rehovot, Israel</b>	CITIZENSHIP <b>Israeli</b>	
POST OFFICE ADDRESS <b>Hanassi Harishon Street 33, 76303 Rehovot, Israel</b>		
FULL NAME OF SECOND JOINT INVENTOR <b>Lior CARMON</b>	INVENTOR'S SIGNATURE	DATE <b>18/1/2001</b>
RESIDENT <b>Tel-Aviv, Israel</b>	CITIZENSHIP <b>Israeli</b> <b>ILX</b>	
POST OFFICE ADDRESS <b>Kehilat-Kiyov Street 6, 69989 Tel-Aviv, Israel</b>		
FULL NAME OF THIRD JOINT INVENTOR <b>Boaz TIROSH</b>	INVENTOR'S SIGNATURE	DATE
RESIDENT <b>Kiryat Ono, Israel</b>	CITIZENSHIP <b>Israeli</b>	
POST OFFICE ADDRESS <b>Stern Street 9, 55602 Kiryat Ono, Israel</b>		
FULL NAME OF FOURTH JOINT INVENTOR <b>Erez BAR-HAIM</b>	INVENTOR'S SIGNATURE	DATE
RESIDENT <b>Yavne, Israel</b>	CITIZENSHIP <b>Israeli</b>	
POST OFFICE ADDRESS <b>Hatsivony Street 25/8, 81573 Yavne, Israel</b>		
FULL NAME OF FIFTH JOINT INVENTOR <b>Adrian PAZ</b>	INVENTOR'S SIGNATURE	DATE
RESIDENT <b>Petach Tikva, Israel</b>	CITIZENSHIP <b>Israeli</b>	
POST OFFICE ADDRESS <b>Wolfson Street 25, 49541 Petach Tikva, Israel</b>		
FULL NAME OF SIXTH JOINT INVENTOR <b>Matityahu FRIDKIN</b>	INVENTOR'S SIGNATURE	DATE
RESIDENT <b>Rehovot, Israel</b>	CITIZENSHIP <b>Israeli</b>	
POST OFFICE ADDRESS <b>Miller Street 23, 76284 Rehovot, Israel</b>		
FULL NAME OF SEVENTH JOINT INVENTOR <b>Cheryl FITZER-ATTAS</b>	INVENTOR'S SIGNATURE	DATE
RESIDENT <b>Rehovot, Israel</b>	CITIZENSHIP <b>US/Israeli</b>	
POST OFFICE ADDRESS <b>Sapir Street 9, 76227 Rehovot, Israel</b>		

ALL INVENTORS MUST REVIEW APPLICATION AND DECLARATION BEFORE SIGNING ALL ALTERATIONS MUST BE INITIALED AND DATED BY ALL INVENTORS PRIOR TO EXECUTION NO ALTERATIONS CAN BE MADE AFTER THE DECLARATION IS SIGNED ALL PAGES OF DECLARATION MUST BE SEEN BY ALL INVENTORS

## Combined Declaration for Patent Application and Power of Attorney

As a below-named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name; and that I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

TUMOR ASSOCIATED ANTIGEN PEPTIDES AND USE OF SAME AS ANTI-TUMOR VACCINES

the specification of which (check one)

- [ ] is attached hereto;  
[ ] was filed in the United States under 35 U.S.C. §111 on \_\_\_\_\_, as  
U.S. Appln. No. \_\_\_\_\_\*; or  
[X] was/will be filed in the U.S. under 35 U.S.C. §371 by entry into the U.S. national stage of an  
international (PCT) application, PCT/IL99/00417; filed 29 July 1999, entry requested on  
\_\_\_\_\_\*; national stage application received U.S. Appln. No. \_\_\_\_\_\*; §371/§102(e)  
date \_\_\_\_\_\* (\* if known)

and was amended on \_\_\_\_\_ (if applicable).  
*(include dates of amendments under PCT Art. 19 and 34 if PCT)*

I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above; and I acknowledge the duty to disclose to the Patent and Trademark Office (PTO) all information known by me to be material to patentability as defined in 37 C.F.R. §1.56.

I hereby claim foreign priority benefits under 35 U.S.C. §§ 119 and 365 of any prior foreign application(s) for patent or inventor's certificate, or prior PCT application(s) designating a country other than the U.S., listed below with the "Yes" box checked and have also identified below any such application having a filing date before that of the application on which priority is claimed:

<u>125608</u>	<u>Israel</u>	<u>30 July 1998</u>	[X]	[ ]
(Number)	(Country)	(Day Month Year Filed)	YES	NO
<u>          </u>	<u>          </u>	<u>          </u>	[ ]	[ ]
(Number)	(Country)	(Day Month Year Filed)	YES	NO

I hereby claim the benefit under 35 U.S.C. §120 of any prior U.S. non-provisional application(s) or prior PCT application(s) designating the U.S. listed below, or under §119(e) of any prior U.S. provisional applications listed below, and, insofar as the subject matter of each of the claims of this application is not disclosed in such U.S. or PCT application in the manner provided by the first paragraph of 35 U.S.C. §112, I acknowledge the duty to disclose to the PTO all information as defined in 37 C.F.R. §1.56(a) which occurred between the filing date of the prior application and the national filing date of this application:

<u>          </u>	<u>          </u>	<u>          </u>
(Application No.)	(Day Month Year Filed)	(Status: patented, pending, abandoned)
<u>          </u>	<u>          </u>	<u>          </u>
(Application No.)	(Day Month Year Filed)	(Status: patented, pending, abandoned)
<u>          </u>	<u>          </u>	<u>          </u>
(Application No.)	(Day Month Year Filed)	(Status: patented, pending, abandoned)

As a named inventor, I hereby appoint the following registered practitioners to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith:

**All of the practitioners associated with Customer Number 001444**

Direct all correspondence to the address associated with **Customer Number 001444**; i.e.,

**BROWDY AND NEIMARK, P.L.L.C.**  
624 Ninth Street, N.W.  
Washington, D.C. 20001-5303  
(202) 628-5197

The undersigned hereby authorizes the U.S. Attorneys or Agents appointed herein to accept and follow instructions from Yeda Research and Development Co. Ltd. as to any action to be taken in the U.S. Patent and Trademark Office regarding this application without direct communication between the U.S. Attorneys or Agents and the undersigned. In the event of a change of the persons from whom instructions may be taken, the U.S. Attorneys or Agents appointed herein will be so notified by the undersigned.

Title: TUMOR ASSOCIATED ANTIGEN PEPTIDES AND USE OF SAME AS ANTI-TUMOR VACCINES

U.S. Application filed \_\_\_\_\_, Serial No. \_\_\_\_\_

PCT Application filed 29 July 1999, Serial No. PCT/IL99/00417

I hereby further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C. §1001 and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

FULL NAME OF FIRST INVENTOR <b>Lea EISENBACH</b>	INVENTOR'S SIGNATURE	DATE
RESIDENT <b>Rehovot, Israel</b>	CITIZENSHIP <b>Israeli</b>	
POST OFFICE ADDRESS <b>Hanassi Harishon Street 33, 76303 Rehovot, Israel</b>		
FULL NAME OF SECOND JOINT INVENTOR <b>Lior CARMON</b>	INVENTOR'S SIGNATURE	DATE
RESIDENT <b>Tel-Aviv, Israel</b>	CITIZENSHIP <b>Israeli</b>	
POST OFFICE ADDRESS <b>Kehilat-Kiyov Street 6, 69989 Tel-Aviv, Israel</b>		
FULL NAME OF THIRD JOINT INVENTOR <b>Boaz TIROSH</b>	INVENTOR'S SIGNATURE	DATE
RESIDENT <b>Kiryat Ono, Israel</b>	CITIZENSHIP <b>Israeli</b>	
POST OFFICE ADDRESS <b>Stern Street 9, 55602 Kiryat Ono, Israel</b>		
FULL NAME OF FOURTH JOINT INVENTOR <b>Erez BAR-HAIM</b>	INVENTOR'S SIGNATURE	DATE
RESIDENT <b>Yavne, Israel</b>	CITIZENSHIP <b>Israeli</b>	
POST OFFICE ADDRESS <b>Hatsivony Street 25/8, 81573 Yavne, Israel</b>		
FULL NAME OF FIFTH JOINT INVENTOR <b>Adrian PAZ</b>	INVENTOR'S SIGNATURE <i>[Signature]</i>	DATE <b>21/1/2001</b>
RESIDENT <b>Petach Tikva, Israel</b>	CITIZENSHIP <b>Israeli</b> <b>ILX</b>	
POST OFFICE ADDRESS <b>Eliezer Friedman Str 13, 49541 Petach Tikva, Israel</b>		
FULL NAME OF SIXTH JOINT INVENTOR <b>Matityahu FRIDKIN</b>	INVENTOR'S SIGNATURE	DATE
RESIDENT <b>Rehovot, Israel</b>	CITIZENSHIP <b>Israeli</b>	
POST OFFICE ADDRESS <b>Miller Street 23, 76284 Rehovot, Israel</b>		
FULL NAME OF SEVENTH JOINT INVENTOR <b>Cheryl FITZER-ATTAS</b>	INVENTOR'S SIGNATURE	DATE
RESIDENT <b>Rehovot, Israel</b>	CITIZENSHIP <b>US/Israeli</b>	
POST OFFICE ADDRESS <b>Sapir Street 9, 76227 Rehovot, Israel</b>		

ALL INVENTORS MUST REVIEW APPLICATION AND DECLARATION BEFORE SIGNING. ALL ALTERATIONS MUST BE INITIALED AND DATED BY ALL INVENTORS PRIOR TO EXECUTION. NO ALTERATIONS CAN BE MADE AFTER THE DECLARATION IS SIGNED. ALL PAGES OF DECLARATION MUST BE SEEN BY ALL INVENTORS.

**Combined Declaration for Patent Application and Power of Attorney**

As a below-named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name; and that I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

TUMOR ASSOCIATED ANTIGEN PEPTIDES AND USE OF SAME AS ANTI-TUMOR VACCINES

the specification of which (check one)

- [ ] is attached hereto;  
 [ ] was filed in the United States under 35 U.S.C. §111 on \_\_\_\_\_, as  
 U.S. Appl. No. \_\_\_\_\_\*; or  
 [X] was/will be filed in the U.S. under 35 U.S.C. §371 by entry into the U.S. national stage of an  
 international (PCT) application, PCT/IL99/00417; filed 29 July 1999, entry requested on  
 \_\_\_\_\_\*; national stage application received U.S. Appl. No. \_\_\_\_\_\*; §371/§102(e)  
 date \_\_\_\_\_\* (\* if known)

and was amended on \_\_\_\_\_ (if applicable).

*(include dates of amendments under PCT Art. 19 and 34 if PCT)*

I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above; and I acknowledge the duty to disclose to the Patent and Trademark Office (PTO) all information known by me to be material to patentability as defined in 37 C.F.R. §1.56.

I hereby claim foreign priority benefits under 35 U.S.C. §§ 119 and 365 of any prior foreign application(s) for patent or inventor's certificate, or prior PCT application(s) designating a country other than the U.S., listed below with the "Yes" box checked and have also identified below any such application having a filing date before that of the application on which priority is claimed:

<u>125608</u>	<u>Israel</u>	<u>30 July 1998</u>	[X]	[ ]
(Number)	(Country)	(Day Month Year Filed)	YES	NO
<u>          </u>	<u>          </u>	<u>          </u>	[ ]	[ ]
(Number)	(Country)	(Day Month Year Filed)	YES	NO

I hereby claim the benefit under 35 U.S.C. §120 of any prior U.S. non-provisional application(s) or prior PCT application(s) designating the U.S. listed below, or under §119(e) of any prior U.S. provisional applications listed below, and, insofar as the subject matter of each of the claims of this application is not disclosed in such U.S. or PCT application in the manner provided by the first paragraph of 35 U.S.C. §112, I acknowledge the duty to disclose to the PTO all information as defined in 37 C.F.R. §1.56(a) which occurred between the filing date of the prior application and the national filing date of this application:

<u>          </u>	<u>          </u>	<u>          </u>
(Application No.)	(Day Month Year Filed)	(Status: patented, pending, abandoned)
<u>          </u>	<u>          </u>	<u>          </u>
(Application No.)	(Day Month Year Filed)	(Status: patented, pending, abandoned)
<u>          </u>	<u>          </u>	<u>          </u>
(Application No.)	(Day Month Year Filed)	(Status: patented, pending, abandoned)

As a named inventor, I hereby appoint the following registered practitioners to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith:

**All of the practitioners associated with Customer Number 001444**

Direct all correspondence to the address associated with Customer Number 001444; i.e.,

**BROWDY AND NEIMARK, P.L.L.C.**  
**624 Ninth Street, N.W.**  
**Washington, D.C. 20001-5303**  
**(202) 628-5197**

The undersigned hereby authorizes the U.S. Attorneys or Agents appointed herein to accept and follow instructions from Yeda Research and Development Co. Ltd. as to any action to be taken in the U.S. Patent and Trademark Office regarding this application without direct communication between the U.S. Attorneys or Agents and the undersigned. In the event of a change of the persons from whom instructions may be taken, the U.S. Attorneys or Agents appointed herein will be so notified by the undersigned.

Title: TUMOR ASSOCIATED ANTIGEN PEPTIDES AND USE OF SAME AS ANTI-TUMOR VACCINES

U.S. Application filed \_\_\_\_\_, Serial No. \_\_\_\_\_

PCT Application filed 29 July 1999, Serial No. PCT/IL99/00417

I hereby further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C. §1001 and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

FULL NAME OF FIRST INVENTOR <b>Lea EISENBACH</b>	INVENTOR'S SIGNATURE	DATE
RESIDENT <b>Rehovot, Israel</b>	CITIZENSHIP <b>Israeli</b>	
POST OFFICE ADDRESS <b>Hanassi Harishon Street 33, 76303 Rehovot, Israel</b>		
FULL NAME OF SECOND JOINT INVENTOR <b>Lior CARMON</b>	INVENTOR'S SIGNATURE	DATE
RESIDENT <b>Tel-Aviv, Israel</b>	CITIZENSHIP <b>Israeli</b>	
POST OFFICE ADDRESS <b>Kehilat-Kiyov Street 6, 69989 Tel-Aviv, Israel</b>		
FULL NAME OF THIRD JOINT INVENTOR <b>Boaz TIROSH</b>	INVENTOR'S SIGNATURE	DATE
RESIDENT <b>Kiryat Ono, Israel</b>	CITIZENSHIP <b>Israeli</b>	
POST OFFICE ADDRESS <b>Stern Street 9, 55602 Kiryat Ono, Israel</b>		
FULL NAME OF FOURTH JOINT INVENTOR <b>Erez BAR-HAIM</b>	INVENTOR'S SIGNATURE	DATE
RESIDENT <b>Yavne, Israel</b>	CITIZENSHIP <b>Israeli</b>	
POST OFFICE ADDRESS <b>Hatsivony Street 25/8, 81573 Yavne, Israel</b>		
FULL NAME OF FIFTH JOINT INVENTOR <b>Adrian PAZ</b>	INVENTOR'S SIGNATURE	DATE
RESIDENT <b>Petach Tikva, Israel</b>	CITIZENSHIP <b>Israeli</b>	
POST OFFICE ADDRESS <b>Wolfson Street 25, 49541 Petach Tikva, Israel</b>		
FULL NAME OF SIXTH JOINT INVENTOR <b>Matityahu FRIDKIN</b>	INVENTOR'S SIGNATURE	DATE
RESIDENT <b>Rehovot, Israel</b>	CITIZENSHIP <b>Israeli</b>	
POST OFFICE ADDRESS <b>Miller Street 23, 76284 Rehovot, Israel</b>		
FULL NAME OF SEVENTH JOINT INVENTOR <b>Cheryl FITZER-ATTAS</b>	INVENTOR'S SIGNATURE <i>Cheryl Fitzer-Attas</i>	DATE <b>1/17/01</b>
RESIDENT <b>Rehovot, Israel</b>	CITIZENSHIP <b>US/Israeli</b> <b>ILX</b>	
POST OFFICE ADDRESS <b>Sapir Street 9, 76227 Rehovot, Israel</b>		

JC07 Rec'd PCT/PTO 30 JAN 2001

## SEQUENCE LISTING

<110> Yeda Research & Development Company Limited at the  
Bio-Technology General Corp.

<120> Tumor Associated Antigen Peptides and Use of Same as  
Anti-Tumor Vaccines

<130> 453-A-PCT

<140>

<141>

<150> 125608

<151> 1998-07-30

<160> 77

<170> PatentIn Ver. 2.1

<210> 1

<211> 9

<212> PRT

<213> Homo sapiens

<400> 1

Phe Leu Leu Val Leu Gly Phe Ile Ile

1

5

<210> 2

<211> 9

<212> PRT

<213> Homo sapiens

<400> 2

Val Leu Pro Ser Val Ala Met Phe Leu

1

5

<210> 3

<211> 9

<212> PRT

<213> Homo sapiens

<400> 3

Leu Val Leu Gly Phe Ile Ile Ala Leu  
1 5

<210> 4  
<211> 9  
<212> PRT  
<213> Homo sapiens

<400> 4  
Lys Val Val Thr Ser Ser Phe Val Val  
1 5

<210> 5  
<211> 9  
<212> PRT  
<213> Homo sapiens

<400> 5  
Leu Val Pro Gly Thr Lys Phe Tyr Ile  
1 5

<210> 6  
<211> 9  
<212> PRT  
<213> Homo sapiens

<400> 6  
Leu Leu Pro Ile Arg Thr Leu Pro Leu  
1 5

<210> 7  
<211> 9  
<212> PRT  
<213> Homo sapiens

<400> 7  
Tyr Leu Val Lys Lys Gly Thr Ala Thr  
1 5

<210> 8  
<211> 9  
<212> PRT  
<213> Homo sapiens

<400> 8  
Ser Leu Phe Ala Glu Thr Ile Trp Val  
1 5

<210> 9  
<211> 9  
<212> PRT  
<213> Homo sapiens

<400> 9  
Met Leu Ile Ala Met Tyr Phe Tyr Thr  
1 5

<210> 10  
<211> 9  
<212> PRT  
<213> Homo sapiens

<400> 10  
Leu Met Trp Thr Leu Pro Val Met Leu  
1 5

<210> 11  
<211> 9  
<212> PRT  
<213> Homo sapiens

<400> 11  
Met Leu Ile Val Tyr Ile Phe Glu Cys  
1 5

<210> 12  
<211> 9  
<212> PRT



<213> Homo sapiens

<400> 12

Tyr Ile Phe Glu Cys Ala Ser Cys Ile  
1 5

<210> 13

<211> 9

<212> PRT

<213> Homo sapiens

<400> 13

Leu Val Leu Met Leu Ile Val Tyr Ile  
1 5

<210> 14

<211> 9

<212> PRT

<213> Homo sapiens

<400> 14

Ala Leu Cys Arg Arg Arg Ser Met Val  
1 5

<210> 15

<211> 9

<212> PRT

<213> Homo sapiens

<400> 15

Leu Leu Ser Gly Leu Ser Leu Phe Ala  
1 5

<210> 16

<211> 9

<212> PRT

<213> Homo sapiens

<400> 16

Phe Leu Leu Val Val Gly Leu Ile Val

1

5

&lt;210&gt; 17

&lt;211&gt; 9

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 17

Leu Val Val Gly Leu Ile Val Ala Leu

1

5

&lt;210&gt; 18

&lt;211&gt; 9

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 18

Lys Val Val Lys Ser Asp Phe Val Val

1

5

&lt;210&gt; 19

&lt;211&gt; 9

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 19

Thr Leu Pro Val Gln Thr Leu Pro Leu

1

5

&lt;210&gt; 20

&lt;211&gt; 9

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 20

Asp Leu His Val Ile Ser Asn Asp Val

1

5

<210> 21

<211> 9

<212> PRT

<213> Homo sapiens

<400> 21

Val Leu Val His Pro Gln Trp Val Leu

1

5

<210> 22

<211> 9

<212> PRT

<213> Homo sapiens

<400> 22

Phe Leu Arg Pro Gly Asp Asp Ser Ser

1

5

<210> 23

<211> 9

<212> PRT

<213> Homo sapiens

<400> 23

Ala Leu Gly Thr Thr Cys Tyr Ala Ser

1

5

<210> 24

<211> 9

<212> PRT

<213> Homo sapiens

<400> 24

Lys Leu Gln Cys Val Asp Leu His Val

1

5

<210> 25

<211> 9

<212> PRT

<213> Homo sapiens

&lt;400&gt; 25

Glu Leu Ala His Tyr Asp Val Leu Leu

1

5

&lt;210&gt; 26

&lt;211&gt; 9

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 26

Asn Leu Asn Gly Ala Gly Asp Pro Leu

1

5

&lt;210&gt; 27

&lt;211&gt; 9

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 27

Thr Leu Arg Val Asp Cys Thr Pro Leu

1

5

&lt;210&gt; 28

&lt;211&gt; 9

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 28

Met Met Asn Asp Gln Leu Met Phe Leu

1

5

&lt;210&gt; 29

&lt;211&gt; 9

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 29

Ala Leu Phe Asp Ile Glu Ser Lys Val

1

5

<210> 30  
<211> 9  
<212> PRT  
<213> Homo sapiens

<400> 30  
Leu Leu His Glu Thr Asp Ser Ala Val  
1 5

<210> 31  
<211> 9  
<212> PRT  
<213> Homo sapiens

<400> 31  
Val Leu Ala Lys Glu Leu Lys Phe Val  
1 5

<210> 32  
<211> 9  
<212> PRT  
<213> Homo sapiens

<400> 32  
Ile Leu Leu Trp Gln Pro Ile Pro Val  
1 5

<210> 33  
<211> 9  
<212> PRT  
<213> Homo sapiens

<400> 33  
Asp Leu Phe Gly Ile Trp Ser Lys Val  
1 5

<210> 34

<211> 9  
<212> PRT  
<213> Homo sapiens

<400> 34  
Pro Leu Glu Arg Phe Ala Glu Leu Val  
1 5

<210> 35  
<211> 9  
<212> PRT  
<213> Homo sapiens

<400> 35  
Lys Gln Gly Asn Phe Asn Ala Trp Val  
1 5

<210> 36  
<211> 9  
<212> PRT  
<213> Homo sapiens

<400> 36  
Asn Leu Leu Arg Arg Met Trp Val Thr  
1 5

<210> 37  
<211> 9  
<212> PRT  
<213> Homo sapiens

<400> 37  
Asn Leu Phe Glu Thr Pro Ile Leu Ala  
1 5

<210> 38  
<211> 9  
<212> PRT  
<213> Homo sapiens

&lt;400&gt; 38

Asn Leu Phe Glu Thr Pro Val Glu Ala

1

5

&lt;210&gt; 39

&lt;211&gt; 9

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 39

Gly Leu Gln His Trp Val Pro Glu Leu

1

5

&lt;210&gt; 40

&lt;211&gt; 9

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 40

Val Gln Phe Val Ala Ser Tyr Lys Val

1

5

&lt;210&gt; 41

&lt;211&gt; 9

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 41

Arg Leu Leu Ala Ala Leu Cys Gly Ala

1

5

&lt;210&gt; 42

&lt;211&gt; 9

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 42

Leu Leu Leu Leu Thr Val Leu Thr Val

1

5

<210> 43  
<211> 9  
<212> PRT  
<213> Homo sapiens

<400> 43  
Leu Leu Leu Thr Val Leu Thr Val Val  
1 5

<210> 44  
<211> 9  
<212> PRT  
<213> Homo sapiens

<400> 44  
Phe Leu Ser Phe His Ile Ser Asn Leu  
1 5

<210> 45  
<211> 9  
<212> PRT  
<213> Homo sapiens

<400> 45  
Leu Leu Val Leu Val Cys Val Leu Val  
1 5

<210> 46  
<211> 9  
<212> PRT  
<213> Homo sapiens

<400> 46  
Ala Leu Leu Val Leu Val Cys Val Leu  
1 5

<210> 47  
<211> 9



<212> PRT

<213> Homo sapiens

<400> 47

Ser Leu Ser Tyr Thr Asn Pro Ala Val

1

5

<210> 48

<211> 9

<212> PRT

<213> Homo sapiens

<400> 48

Asn Leu Thr Ile Ser Asp Val Ser Val

1

5

<210> 49

<211> 9

<212> PRT

<213> Homo sapiens

<400> 49

Ala Leu Ala Ser Thr Ala Pro Pro Val

1

5

<210> 50

<211> 9

<212> PRT

<213> Homo sapiens

<400> 50

Ala Ile Leu Cys Trp Thr Phe Trp Val

1

5

<210> 51

<211> 9

<212> PRT

<213> Homo sapiens

<400> 51

Phe Ile Leu Met Phe Ile Val Tyr Ala  
1 5

<210> 52  
<211> 9  
<212> PRT  
<213> Homo sapiens

<400> 52  
Leu Thr Ala Glu Cys Ile Phe Phe Val  
1 5

<210> 53  
<211> 9  
<212> PRT  
<213> Homo sapiens

<400> 53  
Met Leu Gln Asp Asn Cys Cys Gly Val  
1 5

<210> 54  
<211> 9  
<212> PRT  
<213> Homo sapiens

<400> 54  
Ile Leu Cys Trp Thr Phe Trp Val Leu  
1 5

<210> 55  
<211> 9  
<212> PRT  
<213> Homo sapiens

<400> 55  
Lys Ile Leu Leu Ala Tyr Phe Ile Leu  
1 5

<210> 56  
<211> 9  
<212> PRT  
<213> Homo sapiens

<400> 56  
Phe Val Gly Ile Cys Leu Phe Cys Leu  
1 5

<210> 57  
<211> 9  
<212> PRT  
<213> Homo sapiens

<400> 57  
Val Leu Leu Ser Val Ala Met Phe Leu  
1 5

<210> 58  
<211> 9  
<212> PRT  
<213> Homo sapiens

<400> 58  
Leu Leu Ser Val Ala Met Phe Leu Leu  
1 5

<210> 59  
<211> 9  
<212> PRT  
<213> Homo sapiens

<400> 59  
Ile Leu Gly Ser Leu Pro Phe Phe Leu  
1 5

<210> 60  
<211> 9  
<212> PRT

<213> Homo sapiens

<400> 60

Ile Leu Asn Ala Tyr Leu Val Arg Val  
1 5

<210> 61

<211> 9

<212> PRT

<213> Homo sapiens

<400> 61

Phe Leu Leu Val Gly Phe Ala Gly Ala  
1 5

<210> 62

<211> 9

<212> PRT

<213> Homo sapiens

<400> 62

Asn Leu Gln Pro Gln Leu Ala Ser Val  
1 5

<210> 63

<211> 9

<212> PRT

<213> Homo sapiens

<400> 63

Cys Met Phe Asp Ser Lys Glu Ala Leu  
1 5

<210> 64

<211> 9

<212> PRT

<213> Homo sapiens

<400> 64

Tyr Leu Tyr Val Leu Val Asp Ser Ala

1

5

&lt;210&gt; 65

&lt;211&gt; 9

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 65

Tyr Met Asp Gly Thr Met Ser Gln Val

1

5

&lt;210&gt; 66

&lt;211&gt; 9

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 66

Lys Met Ala Arg Phe Ser Tyr Ser Val

1

5

&lt;210&gt; 67

&lt;211&gt; 9

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 67

Gly Leu Val Met Asp Glu His Leu Val

1

5

&lt;210&gt; 68

&lt;211&gt; 9

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 68

Phe Leu Pro Gly Cys Asp Gly Leu Val

1

5

<210> 69  
<211> 9  
<212> PRT  
<213> Homo sapiens

<400> 69  
Cys Met Leu Gly Ser Phe Cys Ala Cys  
1 5

<210> 70  
<211> 9  
<212> PRT  
<213> Homo sapiens

<400> 70  
Tyr Leu Ala Phe Arg Asp Asp Ser Ile  
1 5

<210> 71  
<211> 9  
<212> PRT  
<213> Homo sapiens

<400> 71  
Trp Leu Pro Lys Lys Cys Ser Leu Cys  
1 5

<210> 72  
<211> 9  
<212> PRT  
<213> Homo sapiens

<400> 72  
Cys Leu Asn Gly Gly Thr Cys Met Leu  
1 5

<210> 73  
<211> 9  
<212> PRT  
<213> Homo sapiens

&lt;400&gt; 73

Met Leu Val Gly Ile Cys Leu Ser Ile

1

5

&lt;210&gt; 74

&lt;211&gt; 9

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 74

Phe Glu Leu Gly Leu Val Ala Gly Leu

1

5

&lt;210&gt; 75

&lt;211&gt; 9

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 75

Lys Met Val Arg Phe Ser Tyr Ser Val

1

5

&lt;210&gt; 76

&lt;211&gt; 9

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 76

Cys Leu Asn Glu Gly Thr Cys Met Leu

1

5

&lt;210&gt; 77

&lt;211&gt; 9

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens


&lt;400&gt; 77

Met Leu Ala Gly Ile Cys Leu Ser Ile

1

5

United States Patent & Trademark Office  
Office of Initial Patent Examination — Scanning Division



Application deficiencies found during scanning:

☐ Page(s) \_\_\_\_\_ of \_\_\_\_\_ were not present  
for scanning. (Document title)

☐ Page(s) \_\_\_\_\_ of \_\_\_\_\_ were not present  
for scanning. (Document title)

\* Scanned copy is best available. *drawings Figure are dark*